

①  **Europäisches Patentamt**
European Patent Office
Office européen des brevets

⑪ Publication number:

0 249 676
A2

⑫

EUROPEAN PATENT APPLICATION

⑰ Application number: 87101137.5

⑮ Int. Cl.: **C 12 N 15/00, C 12 P 21/02,**
C 12 N 1/20, C 12 N 5/00,
A 01 H 1/00

⑱ Date of filing: 28.01.87

⑳ Priority: 28.01.86 DK 412/86

⑦ Applicant: **AKTIESELSKABET DE DANSKE**
SUKKERFABRIKKER, Langebrogade 5,
DK-1411 Copenhagen K (DK)

④③ Date of publication of application: 23.12.87
Bulletin 87/52

⑦② Inventor: **Marcker, Kjeld Adrian, 1, Toftevej, DK-8250 Ege**
(DK)
Inventor: **Jensen, Jens Stougaard, 10, J.M. Morksgade 1.**
sal, DK-8000 Aarhus C (DK)

④④ Designated Contracting States: **AT BE CH DE ES FR GB**
GR IT LI LU NL SE

⑦④ Representative: **Vossius & Partner,**
Siebertstrasse 4 P.O. Box 86 07 67,
D-8000 München 86 (DE)

⑤④ Method for the expression of genes in plants.

⑤⑦ A method for the expression of genes in plants, parts of plants, and plant cell cultures, in which a DNA fragment is used comprising an inducible plant promoter of root nodule-specific genes, DNA-fragments comprising an inducible plant promoter, to be used when carrying out the method, said DNA-fragments being identical with, derived from or comprising a 5' flanking region of root nodule-specific genes of any origin as well as plasmids and transformed *Agrobacterium rhizogenes*-strain which can be used when carrying out the method.

EP 0 249 676 A2

TITLE MODIFIED
see front page

0249676

A method for the expression of genes in plants,
parts of plants, and plant cell cultures, and DNA
fragments, plasmids, and transformed microorganisms
to be used when carrying out the method, as well
5 as the use thereof for the expression of genes in
plants, parts of plants, and plant cell cultures.

The invention relates to a novel method for the
expression of genes in plants, parts of plants,
and plant cell cultures, as well as DNA fragments
10 and plasmids comprising said DNA fragments to be
used when carrying out the method. The invention
furthermore relates to transformed plants, parts
of plants and plant cells.

The invention relates to this method for the ex-
15 pression of genes of any origin under control of
an inducible, root nodule specific promoter.

The invention relates especially to this method
for the expression of root nodule-specific genes
in transformed plants including both leguminous
20 plants and other plants.

The invention relates furthermore to DNA fragments
comprising an inducible plant promoter to be used
when carrying out the method, as well as plasmids
comprising said DNA fragments.

25 In the specification i.a. the following terms are
used:

Root nodule-specific genes: Plant genes active
only in the root nodules of leguminous plants, or

genes with an increased expression in root nodules. Root nodule-specific plant genes are expressed at predetermined stages of development and are activated in a coordinated manner during the symbiosis
5 whereby a nitrogen fixation takes place and the fixed nitrogen is utilized in the metabolism of the plant.

Inducible plant promoter: Generally is meant a promoter-active 5' flanking region from plant genes
10 inducible from a low activity to a high activity. In relation to the present invention "inducible plant promoter" means a promoter derived from, contained in or being identical with a 5' flanking region including a leader sequence of root nodule-
15 specific genes and being capable of promoting and regulating the expression of a gene as characterised in relation to the present invention.

Leader sequence: Generally is meant a DNA sequence being transcribed into a mRNA, but not further
20 translated into protein. The leader sequence comprises thus the DNA fragment from the start of the transcription to the ATG codon constituting the start of the translation. In relation to the present invention "leader sequence" means a short DNA frag-
25 ment contained in the above inducible plant promoter and typically comprising 40-70 bp and which may comprise sequences being targets for a posttranscriptional regulation.

Promoter region: A DNA fragment containing a pro-
30 moter which comprises target sequences for RNA polymerase as well as possible activation regions

comprising target sequences for transcriptional effector substances. In the present invention, target sequences for transcriptional effectors may also be situated 3' to the promoter, i.e. in the 5 coding sequences, the intervening sequences or on the 3' flanking region of a root nodule-specific gene.

Furthermore a number of molecular-biological terms generally known to persons skilled in the art are
10 used, including the terms stated below:

CAP (addition) site: The nucleotide of the 5' end of the transcript where 7-methylGTP is added; In the Figures often given also as an asterisk *-marked nucleotide on a given nucleotide sequence.

15 DNA sequence or DNA segment: A linear array of nucleotides interconnected through phosphodiester bonds between the 3' and 5' carbon atoms of adjacent pentoses.

Expression: The process undergone by a structural
20 gene to produce a polypeptide. It is a combination of transcription and translation as well as possible posttranslational modifications.

Flanking regions: DNA sequences surrounding coding regions. 5' flanking regions contain a promoter.
25 3' flanking regions may contain a transcriptional terminator etc.

Gene: A DNA sequence composed of three or four parts, viz. (1) the coding sequence for the gene

product, (2) the sequences in the promoter region which control whether or not the gene will be expressed, (3) those sequences in the 3' end conditioning the transcriptional termination and optionally polyadenylation, as well as (4) intervening sequences, if any.

Intervening sequences: DNA sequences within a gene which are not coding for any peptide fragment. The intervening sequences are transcribed into pre-mRNA and are eliminated by modification of pre-mRNA into mRNA. They are also called introns.

Chimeric gene: A gene composed of parts from various genes. E.g. the chimeric Lbc₃-5'-3'-CAT is composed of a chloroamphenicolacetyltransferase-coding sequence deriving from E. coli and 5' and 3' flanking regulatory regions of the Lbc₃ gene of soybean.

Cloning: The process of obtaining a population of organisms or DNA sequences deriving from one such organism or sequence by asexual reproduction, or more particular a process of isolating a particular organism or part thereof, and the propagation of this subfraction as a homogeneous population.

Coding sequences: DNA sequences determining the amino acid sequence of a polypeptide.

25 Cross-inoculation group: A group of leguminous plant species capable of producing functionally active root nodules with Rhizobium bacteria isolated from root nodules of other species of the group.

Leghemoglobin (Lb): An oxygen-binding protein exclusively synthesized in root nodules. The Lb proteins regulate the oxygen partial pressure in the root nodule tissue and transport oxygen to the bacteroides. In this manner the oxygen-sensitive nitrogenase enzyme is protected. The Lb genes are root nodule-specific genes.

Messenger-RNA (mRNA): RNA molecule produced by transcription of a gene and possibly modification of mRNA. The mRNA molecule mediates the genetic message determining the amino acid sequence of a polypeptide by part of the mRNA molecule being translated into said peptide.

Downstream: A position in a DNA sequence. It is defined relative to the transcriptional direction 5'- 3' of the gene relative to which the position is stated. The 3' flanking region is thus positioned downstream of the gene.

Nucleotide: A monomeric unit of DNA or RNA consisting of a sugar moiety (pentose), a phosphate, and a nitrogenous heterocyclic base. The base is linked to the sugar moiety via a glycosidic bond (1' carbon of the pentose), and this combination of base and sugar is a nucleoside. The base characterises the nucleotide. The four DNA bases are adenine (A), guanine (G), cytosine (C), and thymine (T). The four RNA bases are A, G, C, and uracil (U).

Upstream: A position in a DNA sequence. It is defined relative to the transcriptional direction 5'- 3' of the gene relative to which the position

is stated. The 5' flanking region is thus positioned upstream of this gene.

Plant transformation: Processes leading to incorporation of genes in the genome of plant cells in such a manner that these genes are reliably inherited through mitosis and meiosis or in such a manner that these genes are only maintained for short periods.

Plasmid: An extra-chromosomal double-stranded DNA sequence comprising an intact replicon such that the plasmid is replicated in a host cell. When the plasmid is placed within a unicellular organism, the characteristics of that organism are changed or transformed as a result of the DNA of the plasmid. For instance a plasmid carrying the gene for tetracycline resistance (Tc^R) transforms a cell previously sensitive to tetracycline into one which is resistant to it. A cell transformed by a plasmid is called a transformant.

Polypeptide: A linear array of amino acids interconnected by means of peptide bonds between the α -amino and carboxy groups of adjacent amino acids.

Recombination: The creation of a new DNA molecule by combining DNA fragments of different origin.

Homologous recombination: A recombination between sequences showing a high degree of homology.

Replication: A process reproducing DNA molecules.

Replicon: A self-replicating genetic element possessing an origin for the initiation of DNA replication and genes specifying the functions necessary for a control and a replication thereof.

- 5 Restriction fragment: A DNA fragment resulting from double-stranded cleavage by an enzyme recognizing a specific target DNA sequence.

RNA polymerase: Enzyme effecting the transcription of DNA into RNA.

- 10 Root nodule: Specialized tissue resulting from infection of mainly roots of leguminous plants with Rhizobium bacteria. The tissue is produced by the host plant and comprises therefore plant cells whereas the Rhizobium bacteria upon infection are
15 surrounded by a plant cell membrane and differentiate into bacteroides. Root nodules are produced on other species of plants upon infection of nitrogen-fixing bacteria not belonging to the Rhizobium genus. Root nodule-specific plant genes are also
20 expressed in these nodules.

Southern-hybridization: Denatured DNA is transferred upon size separation in agarose gel to a nitrocellulose membrane. Transferred DNA is analysed for a predetermined DNA sequence or a predetermined
25 gene by hybridization. This process allows a binding of single-stranded, radioactively marked DNA sequences (probes) to complementary single-stranded DNA sequences bound on the membrane. The position of DNA fragments on the membrane binding the probe
30 can subsequently be detected on an X-ray film.

Symbiotic nitrogen fixation: The relationship whereby bacteroides of root nodules convert the nitrogen (dinitrogen) of the air into ammonium utilized by the plant while the plant provides the bacteroides with carbon compounds as a carbon source.

Symbiont: One part of a symbiotic relationship, and especially Rhizobium is called the microsymbiont.

Transformation: The process whereby a cell is incorporating a DNA molecule.

10 Translation: The process of producing a polypeptide from mRNA or:

the process whereby the genetic information present in a mRNA molecule directs the order of specific amino acids during the synthesis of a polypeptide.

15 Transcription: The method of synthesizing a complementary RNA sequence from a DNA sequence.

Vector: A plasmid, phage DNA or other DNA sequences capable of replication in a host cell and having one or a small number of endonuclease recognition sites at which such DNA sequences may be cleaved in a determinable manner without loss of an essential biological function.

25 Traditional plant breeding is based on repeated crossbreeding of plant lines individually carrying desired qualities. The identification of progeny lines carrying all the desired qualities is a particularly time-consuming process as the biochemical

and genetic basis of the qualities is usually unknown. New lines are therefore chosen according to their phenotype, usually after a screening of many lines in field experiments.

- 5 Through the ages a direct connection has existed
between the state of nutrition, i.e. the health,
of the population and the agricultural possibility
of ensuring a sufficient supply of assimilable
nitrogen in order to obtain satisfactory yields.
- 10 Already in the seventeenth century it was discovered
that plants of the family leguminosae including
beyond peas also beans, lupins, soybean, bird's-foot
trefoil, vetches, alfalfa, sainfoin, and trefoil had
an ability of improving crops grown on the habitat
- 15 of these plants. Today it is known that the latter
is due to the fact that the members of the plants
of the family leguminosae are able to produce nitro-
gen reserves themselves. On the roots they carry
bacteria with which they live in symbiosis.
- 20 An infection of the roots of these leguminous plants
with Rhizobium bacteria causes a formation of root
nodules able to convert atmospheric nitrogen into
bound nitrogen, which is a process called nitrogen
fixation.
- 25 Atmospheric nitrogen is thereby converted into forms
which can be utilized by the host plant as well as
by the plants later on growing on the same habitat.

In the nineteenth century the above possibility was
utilized for the supply of nitrogen in order to
30 achieve a novel increase of the crop yield.

The later further increases in the yield have, however, especially been obtained by means of natural fertilizers and nitrogen-containing synthetic fertilizers. The resulting pollution of the environment makes it desirable to provide alternative possibilities of ensuring the supply of nitrogen necessary for the best possible yields obtainable.

It would thus be valuable to make an improvement possible of the existing nitrogen fixation systems in leguminous plants as well as to allow an incorporation of nitrogen fixation systems in other plants.

The recombinant DNA technique and the plant transformation systems developed render it now possible to provide plants with new qualities in a well-controlled manner. These characteristics can derive from not only the same plant species, but also from all other prokaryotic or eukaryotic organisms. The DNA techniques allow further a quick and specific identification of progeny lines carrying the desired qualities. In this manner a specific plant line can be provided with one or more desired qualities in a quick and well-defined manner.

Correspondingly, plant cells can be provided with well defined qualities and subsequently be maintained as plant cell lines by means of known tissue culture methods. Such plant cells can be utilized for the production of chemical and biological products of particular interest such as dyes, flavours, aroma components, plant hormones, pharmaceutical

products, primary and secondary metabolites as well as polypeptides (enzymes).

A range of factors and functions necessary for biological production of a predetermined gene product are known. Both the initiation and regulation of transcription as well as the initiation and regulation of posttranscriptional processes can be characterised.

At the gene level it is known that these functions are mainly carried out by 5' flanking regions. A wide range of 5' flanking regions from prokaryotic and eukaryotic genes has been sequenced, and in view inter alia thereof a comprehensive knowledge has been provided of the regulation of gene expression and of the sub-regions and sequences being of importance for the regulation of expression of the gene. Great differences exist in the regulatory mechanism of prokaryotic and eukaryotic organisms, but many common features apply to the two groups.

The regulation of the expression of gene may take place on the transcriptional level and is then preferably exerted by regulating the initiation frequency of transcription. The latter is well-known and described inter alia by Benjamin Lewin, Gene Expression, John Wiley & Sons, vol. I, 1974, vol. II, Second Edition 1980, vol. III, 1977. As an alternative the regulation may be exerted at the posttranscriptional level, e.g. by the regulation of the frequency of the translation initiation, at the rate of the translation, and of the termination of the translation.

The present invention is based on the surprising finding that 5' flanking regions of root nodule-specific genes, exemplified by the 5' flanking region of the soybean leghemoglobin Lbc₃ gene, can
5 be used for inducible expression of a foreign gene in an alien leguminous plant. The induction and regulation of the promoter is preferably carried out in the form of a regulation and induction at the transscriptonal level and differs thereby
10 from the inducability stated in Patent Application No. 86114704.9, the latter inducability preferably being carried out at the translation level.

The transscripton of both the Lbc₃ gene of the soybean and of a chimeric Lbc₃ gene transferred to
15 bird's-foot trefoil starts at a low level immediately upon the appearance of the root nodules on the plant roots. Subsequently, a high increase of the transscripton takes place immediately before the root nodules turn red. The transscripton of a range
20 of other root nodule-specific genes is initiated exactly at this time. The simultaneous induction of the transscripton of the Lb genes and other root nodule-specific genes means that a common DNA sequence(s) must be present for the various genes
25 controlling this pattern of expression. Thus the leghemoglobin-c₃ gene is a representative of one class of genes and the promoter and the leader sequence, target areas for activation as well as the control elements of the organ specificity of
30 the Lbc₃ gene are representatives of the control elements of a complete gene class.

The promoter of the 5' flanking regions of the Lb genes functions in soybeans and is responsible for the transcription of the Lb genes in root nodules. It is furthermore known, that the efficiency of both the transcription initiation and the subsequent translation initiation on the leader sequence of the Lb genes is high as the Lb proteins constitute approximately 20% of the total protein content in root nodules.

10 The sequence of 5' flanking regions of the four soybean leghemoglobin genes Lba, Lbc₁, Lbc₂, and Lbc₃ appears from the enclosed sequence scheme, scheme 1, wherein the sequences are stated in such a manner that the homology between the four 5' flanking regions appears clearly.

In the sequence scheme "-" indicates that no base is present in the position in question. The names of the genes and the base position counted upstream from the ATG start codon are indicated to the right of the sequence scheme. Furthermore the important sequences have been underlined.

As it appears from the sequence scheme a distinct degree of homology exists between the four 5' flanking regions, and in the position 23-24 bp upstream from the CAP addition site they all contain a TATATAAA sequence corresponding to the "TATA" box which in eukaryotic cells usually are located a corresponding number of bp upstream from the CAP addition site. Furthermore a CCAAG sequence is present 64-72 bp upstream from the CAP addition site, said sequence corresponding to the "CCAAT"

box usually located 70-90 bp upstream from the CAP addition site. From the CAP addition site to the translation start codon, ATG, leader sequences of 52-59 bp are present and show a distinct degree of 5 homology of approx. 75-80%.

In accordance with the present invention it has furthermore been proved, exemplified by the Lbc₃ gene, that the 5' flanking regions of the soybean leghemoglobin genes are functionally active in 10 other plant species. The latter has been proved by fusing the E. coli chloroamphenicol acetyl transferase (CAT) gene with the 5' and 3' flanking regions of the soybean Lbc₃ gene in such a manner that the expression of the CAT gene is controlled 15 by the Lb promoter. This fusion fragment was cloned into the integration vectors pAR1 and pAR22, whereby the plasmids pAR29 and pAR30 were produced. Through homologous recombination the latter plasmids were integrated into the Agrobacterium rhizogenes 20 T DNA region. The transformation of Lotus corniculatus (bird's-foot trefoil) plants, i.e. transfer of the T DNA region, was obtained by wound infection on the hypocotyl. Roots developed from the transformed plant cells were cultivated in vitro and 25 freed from A. rhizogenes bacteria by means of antibiotics. Completely regenerated plants were produced by these root cultures in a conventional manner through somatic embryogenesis or organogenesis.

Regenerated plants were subsequently inoculated 30 with Rhizobium loti bacteria and root nodules for analysis were harvested. Transcription and translation of the chimeric Lbc₃ CAT gene could subse-

quently be detected in root nodules on transformed plants as the activity of the produced chloroamphenicol acetyl transferase enzyme.

The conclusion can subsequently be made that the 5 promoter-containing 5' flanking regions of root nodule-specific genes exemplified by the soybean Lbc₃ promoter are functionally active in foreign plants. The latter is a surprising observation as root nodules are only developed as a consequence
10 of a very specific interaction between the leguminous plant and its corresponding Rhizobium micro-symbiont.

Soybeans produce nodules only upon infection by the species Rhizobium japonicum and Lotus corniculatus
15 only upon infection by the species Rhizobium loti. Soybean and Lotus corniculatus belong therefore to two different cross-inoculation groups, each group producing root nodules by means of two different Rhizobium species. The expression of a chimeric
20 soybean gene in Lotus corniculatus proves therefore an unexpected universal regulatory system applying to the expression of root nodule-specific genes. The regulatory DNA sequences involved can be placed on the 5' and 3' flanking regions of the genes,
25 here exemplified by the 2.0 Kb 5' and 0.9 Kb 3' flanking regions of the Lbc₃ gene. This surprising observation allows the use of root nodule-specific promoters and regulatory sequences in any other plant species and any other plant cell line.

30 In other experiments the 5' flanking region of the nodule-specific N23 gene was fused to the CAT gene

and the Lbc₃ 3' flanking region in such a manner that the expression of the CAT gene is controlled by the N23 promoter. This fusion fragment was cloned into the integration vector pAR22 producing the plasmid N23-CAT which was subsequently recombined into A. rhizogenes and transferred to Lotus corniculatus and Trifolium repens (white clover) by the previously described method. The root nodule-specific expression of the transferred N23-CAT gene
10 obtained in L. corniculatus infected with Rhizobium loti and in T. repens infected with Rhizobium trifolii further demonstrated that expression of root nodule-specific genes is independent of the plant species and Rhizobium species. A universal regulatory system therefore regulates the expression
15 of root nodule-specific genes in the different symbiotic systems formed between legumes and the Rhizobium species of the various cross-inoculation groups.

20 It is known from European Patent Application EP 122,791.A1 that plant genes from one species, by Agrobacterium mediated transformation, can be transferred into a different plant species. It is also known from EP 122,791.A1 that a transferred gene
25 encoding the seed storage protein "Phaseolin" can be expressed into tobacco and alfalfa. From the literature it is also known that this expression is seed specific (Sengupta-Gopalan et al. 1985, Proc. Natl. Acad. Sci. 82, 33203324).

30 The present invention therefore relates to a novel method for the expression of transferred genes in a root nodule-specific manner, using DNA regulatory

sequences from the 5' promoter region, the coding region, or the 3' flanking region of root nodule-specific genes, here exemplified by the leghemoglobin Lbc₃ gene and the N23 gene. This method is
5 distinct from both the method of Agrobacterium mediated transformation and expression of the seed storage protein phaseolin gene characterised in EP 122,791.A1. Expression of the transferred phaseolin gene in EP 122,791.A1 only demonstrates that the
10 phaseolin gene family with its particular regulatory requirements can be expressed in tobacco and alfalfa. It does not demonstrate nor predict that any other genes with their particular regulatory requirements can be expressed in any other plants or
15 plant tissue.

An object of the present invention is to provide a possibility of expressing desired genes in plants, parts of plants, and plant cell cultures.

A further object of the invention is to render it
20 possible to express genes of any origin by the control of an inducible root nodule-specific promoter.

A particular object of the invention is to provide a possibility of expressing desired genes in legu-
25 minous plants.

A still further particular object of the invention is to provide a possibility of expressing root nodule-specific genes in non-leguminous plants.

Further objects of the invention are to improve the

existing nitrogen-fixing systems in leguminous plants as well as to incorporate nitrogen-fixing systems in other plants.

A further object of the invention is to provide a possibility of in certain cases allowing the use of specific sequences of the 3' flanking region, of the coding sequence, and of intervening sequences to influence the regulation of the root nodule-specific promoter.

10 Furthermore it is an object of the invention to provide plasmids comprising the above mentioned inducible plant promoter.

Further objects of the invention appear immediately from the following description.

15 The method according to the invention for the expression of genes in plants, parts of plants, and plant cell cultures is carried out by introducing into a cell thereof a recombinant DNA segment containing both the gene to be expressed and a 5' flanking region comprising a promoter sequence, and
20 optionally a 3' flanking region, and culturing of the transformed cells in a growth medium, said method being characterised by using as the recombinant DNA segment a DNA fragment comprising an
25 inducible plant promoter (as defined) from root nodule-specific genes. If desired the transformed cells are regenerated to plants.

The method according to the invention allows in a well defined manner an expression of foreign genes

in plants, parts of plants, and plant cell cultures, in this connection especially genes providing the plants with desired properties such as for instance a resistance to plant diseases and increased content
5 of valuable polypeptides.

A further use is the preparation of valuable products such as for instance dyes, flavourings, plant hormones, pharmaceutical products, primary and secondary metabolites, and polypeptides by means
10 of the method according to the invention in plant cell cultures and plants.

By using the method according to the invention for the expression of root nodule-specific genes it is possible to express root nodule-specific genes
15 necessary for the formation of an active nitrogen-fixing system both in leguminous plants and other plants. The correct developmental control, cf. Example 8, allows the establishment of a symbiotic nitrogen-fixing system in non-leguminous plants. In
20 this manner it is surprisingly possible to improve the existing nitrogen-fixing systems in leguminous plants as well as to incorporate nitrogen-fixing systems in other plants.

The use of the method according to the invention
25 for the expression of foreign genes in root nodules renders it possible to provide leguminous plants with improved properties such as resistance to herbicides and resistance to diseases and pest.

According to a particular embodiment of the method
30 according to the invention a DNA fragment is used

which comprises an inducible plant promoter and which is identical with, derived from, or comprises 5' flanking regions of leghemoglobins genes. In this manner the expression of any gene is obtained.

5 Examples of such DNA fragments are DNA fragments of the four 5' flanking regions of the soybean leghemoglobin genes, viz.

Lba with the sequence:

```

10 GAGATACATT ATAATAATCT CTCTAGTGTC TATTTATTAT TTTATCTGGT
   GATATATACC TTCTCGTATA CTGTTATTTT TTCAATCTTG TAGATTTACT
   TCTTTTATTT TTATAAAAAA GACTTTATTT TTTTAAAAAA AATAAAGTGA
   ATTTTGAAAA CATGCTCTTT GACAATTTTC TGTTTCCTTT TTCATCATTG
   GGTAAATCT CATAGTGCCT CTATTCAATA ATTTGGGCTC AATTTAATTA
   GTAGAGTCTA CATAAAATTT ACCTTAATAG TAGAGAATAG AGAGTCTTGG
   AAAGTTGGTT TTTCTCGAGG AAGAAAGGAA ATGTTAAAAA CTGTGATATT
15 TTTTTTTTGG ATTAATAGTT ATGTTTATAT GAAAACTGAA AATAAATAAA
   CTAACCATAT TAAATTTAGA ACAACACTTC AATTATTTTT TTAATTTGAT
   TAATTAAAAA ATTATTTGAT TAAATTTTTT AAAAGATCGT TGTTTCTTCT
   TCATCATGCT GATTGACACC CTCCACAAGC CAAGAGAAAC ACATAAGCTT
   TGGTTTTCTC ACTCTCCAAG CCCTCTATAT AAACAAATAT TGGAGTGAAG
   TTGTTGCATA ACTTGCATCG AACAATTAAT AGAAATAACA GAAAATTAAA
   AAAGAAATAT G,

```

20 Lbc₁ with the sequence:

```

TTCTCTTAAT ACAATGGAGT TTTTGTTGAA CATACATACA TTTAAAAAAA
AATCTCTAGT GTCTATTTAC CCGGTGAGAA GCCTTCTCGT GTTTTACACA
CTTTAATATT ATTATATCCT CAACCCACAC AAAAAGAATA CTGTTATATC
TTTCCAAACC TGTAGATTTA TTTATTTATT TATTTATTTT TACAAAGGAG
ACTTCAGAAA AGTAATTACA TAAAGATAGT GAACATCATT TTATTTATTA
25 TAATAAACTT TAAAATCAAA CTTTTTTATA TTTTTTGTTA CCCTTTTCAT
   TATTGGGTGA AATCTCATAG TGAAGCCATT AAATAATTTG GGCTCAAGTT
   TTATTAGTAA AGTCTGCATG AAATTTAACT TAACAATAGA GAGAGTTTTC
   GAAAGGGAGC GAATGTTAAA AAGTGTGATA TTATATTTTA TTTCGATTAA
   TAATTATGTT TACATGAAAA CATACAAAAA AATACTTTTA AATTCAGAAT
   AATACTTAAA ATATTTATTT GCTTAATTGA TTAAGTGAAA ATTATTTGAT
   TAGGATTTTG AAAAGATCAT TGGCTCTTCG TCATGCCGAT TGACACCCCTC
   CACAAGCCAA GAGAAACTTA AGTTGTAAAC TTTCTCACTC CAAGCCTTCT
   ATATAAACAT GTATTGGATG TGAAGTTATT GCATAACTTG CATTGAACAA
30 TAGAAAAATA CAAAAAAAG TAAAAAAGTA GAAAAGAAAT ATG,

```

Lbc₂ with the sequence:

```

TCGAGTTTTT ACTGAACATA CATTATTAA AAAAAACTCT CTAGTGTCCTA
TTTATTCGGC GAGAAGCCTT CTCGTGCTTT ACACACTTTA ATATTATTAT
ATCCCCACCC CCACCAAAAA AAAAAAACT GTTATATCTT TCCAGTACAT
TTATTTCTTA TTTTACAAA GGAAACTTCA CGAAAGTAAT TACAAAAAAG
5 ATAGTGAACA TCATTTTTTT AGTTAAGATG AATTTTAAAA TCACACTTTT
TTATATTTTT TTGTTACCCCT TTTCATTATT GGGTGAAATC TCATAGTGAA
ACTATTAAAT AGTTTGGGCT CAAGTTTTAT TAGTAAAGTC TGCATGAAAT
TTAACTTAAT AATAGAGAGA GTTTTGGAAG GGTAACGAAT GTTAGAAAGT
GTGATATTAT TATAGTTTTA TTTAGATTAA TAATTATGTT TACATGAAAA
TTGACAATTT ATTTTAAAA TTCAGAGTAA TACTTAAATT ACTTATTTAC
TTTAAGATTT TGAAAAGATC ATTTGGCTCT TCATCATGCC GATTGACACC
CTCCACAAGC CAAGAGAAAC TTAAGTTGTA ATTTTCTAA CTCCAAGCCT
10 TCTATATAAA CACGTATTGG ATGTGAAGTT GTTGCAATAC TTGCATTGAA
CAATAGAAAT AACAACAAAG AAAATAAGTG AAAAAAGAAA TATG,

```

and Lbc₃ with the sequence:

```

TATGAAGATT AAAAAATACA CTCATATATA TGCCATAAGA ACCAACAAAA
GACTATTTA AGAAAAGAAA AAAAAACCT GCTACATAAT TTCCAATCTT
15 GTAGATTTAT TTCTTTTATT TTTATAAAGG AGAGTTAAAA AAATTACAAA
ATAAAAATAG TGAACATCGT CTAAGCATTT TTATATAAGA TGAATTTTAA
AAATATAATT TTTTGTCTA AATCGTATGT ATCTTGTCTT AGAGCCATTT
TTGTTTAAAT TGGATAAGAT CACACTATAA AGTTCTTCCT CCGAGTTTGA
TATAAAAAAA ATTGTTTCCC TTTTGATTAT TGGATAAAAT CTCGTAGTGA
CATTATATTA AAAAAATTAG GGCTCAATTT TTATTAGTAT AGTTTGCATA
20 AATTTTAACT TAAAAATAGA GAAAATCTGG AAAAGGGACT GTTAAAAAGT
GTGATATTAG AAATTTGTCT GATATATTAA TATTTTATTT TATATGGAAA
CTAAAAAAT ATATATTAAA ATTTTAAATT CAGAATAATA CTTAAATTAT
TTATTTACTG AAAATGAGTT GATTTAAGTT TTTGAAAAGA TGATTGTCCTC
TTCACCATAC CAATTGATCA CCCTCCTCCA ACAAGCCAAG AGAGACATAA
GTTTTATTAG TTATTCTGAT CACTCTTCAA GCCTTCTATA TAAATAAGTA
TTGGATGTGA AGTTGTTGCA TAACTTGCAT TGAACAATTA ATAGAAATAA
25 CAGAAAAGTA GAAAAGAAAT ATG,

```

A further embodiment of the method according to the invention uses a DNA fragment identical with, derived from or comprising 5' flanking regions of the Lbc₃-5'-3'-CAT gene with the sequence:

```

30 TATGAAGATT AAAAAATACA CTCATATATA TGCCATAAGA ACCAACAAAA
GACTATTTA AGAAAAGAAA AAAAAACCT GCTACATAAT TTCCAATCTT
GTAGATTTAT TTCTTTTATT TTTATAAAGG AGAGTTAAAA AAATTACAAA
ATAAAAATAG TGAACATCGT CTAAGCATTT TTATATAAGA TGAATTTTAA
AAATATAATT TTTTGTCTA AATCGTATGT ATCTTGTCTT AGAGCCATTT
TTGTTTAAAT TGGATAAGAT CACACTATAA AGTTCTTCCT CCGAGTTTGA

```

TATAAAAAAA ATTGTTTCCC TTTTGATTAT TGGATAAAAT CTCGTAGTGA
 CATTATATTA AAAAAATTAG GGCTCAATTT TTATTAGTAT AGTTTGCATA
 AATTTTAACT TAAAAATAGA GAAAATCTGG AAAAGGGGACT GTTAAAAAGT
 GTGATATTAG AAATTGTCG GATATATTAA TATTTTATTT TATATGGAAA
 CTAAAAAAT ATATATTAA ATTTTAAATT CAGATAATA CTTAAATTAT
 5 TTATTACTG AAAATGAGTT GATTTAAGTT TTTGAAAAGA TGATTGTCTC
 TTCACCATAC CAATTGATCA CCCCCTCCA ACAGCCCAAG AGAGACATAA
 GTTTTATTAG TTATTCTGAT CACTCTTCAA GCCTTCTATA TAAATAAGTA
 TTGGATGTGA AGTTGTTGCA TAACCTGCAT TGAACAATTA ATAGAAATAA
 CAGAAAAGTA GAATTCTAAA ATG

- 10 A still further preferred embodiment of the method according to the invention uses a DNA fragment identical with, derived from or comprising 5' flanking regions of the N23 gene with the sequence

10 20 30 40 50 60 70
 GAATTCGAGCTCGCCCGGGGATCGATCCTCTAGAGTCGACCTGCAGCCCAAGCTTGGATCAATCAATTAA
 EcoRI SalI
 80 90 100 110 120 130 140
 15 TTCTATTGAGACACGATTTGAACAATTTTACATTATGAGACTATTTTGGTTTTTTATTGATCCAAA
 150 160 170 180 190 200 210
 AAATTTAAAGCTTTAGATGATGATGAATTGAANNAATATTGTATTAAATNTGAAAAGTTNNNNNGGTTTA
 220 230 240 250 260 270 280
 ATGAATGCTATGATATTGATGGTCTTGATNTATTNNCAGAATTGAAAGTATTAAGAGAAGTGTTAAGAAA
 290 300 310 320 330 340 350
 AGAAGTTAGCACACCAATAGAAGTATTGAGTTATATTAAACTTTAGATTCTTTTCAAATGTTTACATTG
 360 370 380 390 400 410 420
 CATATAGAATTTTATTGACAATCCTTATAACAGTTGCTACTGTTGAAAGACGTTCTTCAAATTAATTA
 430 440 450 460 470 480 490
 20 ACTTAAATCATATCTAAATCAACAATGTTACAAGATAGATTGAATGAGTTAGTTATTTTATCTATTGAA
 500 510 520 530 540 550 560
 AGTAAAGTGTTAGAATTGTTTGATTATAAACTCTGATAAATGATTTTGCAGTTAAAAAACTAGAAGAT
 570 580 590 600 610 620 630
 TAATATAAAATGATATTTTATATAATATATTAAGTCTCTTTAAATTCCTTGTAAGAAAAGACATTTTT
 640 650 660 670 680 690 700
 AAATAATAAAATAAGCAACTCTTAATTTTAAATGAACATCCCTTGTAAACCGAATCTTCCATAATGT
 710 720 730 740 750 760 770
 AAAAATTAATGCTTGATGGAAGTTTTTAATTTGTTCTACTCAATACTCAAAGGGTTGTAATATTTTTTT
 780 790 800 810 820 830 840
 25 TATCATTTTATGTTGTAAATATGAATGCACTAGTAATTAGTTTAAATGATAAAATATATTCTACAGATAT

850 860 870 880 890 900 910
 ATTTCTGTCTCTTGGCAACTCGTGAGAATTGAATATATTATAAAGATGAAAGGTCGTTACAATTTTTTTT
 920 930 940 950 960 970 980
 AGAATAAATATTTATATACAATTCCTAGATTTTGTATATAAATTACATATTGTATGAGTATAAATACAT
 990 1000 1010 1020 1030 1040 1050
 GAGCACACACCAAACTAGTCTCAAATTAAGTAAGGTGCTAATTATTAGCGGCTAGCTAAGTAACCAAGTA
 DdeI
 ATTAATG

5 In a particularly preferred embodiment of the method
 according to the invention a 3' flanking region of
 root nodule-specific genes is furthermore used, in
 particular sequences of the 3' flanking region ca-
 pable of influencing the activity or regulation of
 10 a promotor of the root nodule-specific genes or
 the transcription termination, or capable of in-
 fluencing the yield of the desired gene product in
 another manner.

Examples of such 3' flanking regions are the four
 15 3' flanking regions of the soybean leghemoglobin
 genes, viz.

Lba with the sequence:

1590 1620
 TAA TTA GTA TCT ATT GCA GTA AAG TGT AAT AAA TAA ATC TTG
 1650 1680
 TTT CAC TAT AAA ACT TGT TAC TAT TAG ACA AGG GCC TGA TAC AAA ATG TTG GTT AAA ATA
 1710 1740
 20 ATG GAA TTA TAT AGT ATT GGA TAA AAA TCT TAA GGT TAA TAT TCT ATA TTT GCG TAG GTT
 1770 1800
 TAT GCT TGT GAA TCA TTA TCG GTA TTT TTT TTC CTT TCT GAT AAT TAA TCG GTA AAT TA
 1830 1860
 ACA AAT AAG TTC AAA ATG ATT TAT ATG TTT CAA AAT TAT TTT AAC AGC AGG TAA AAT GTT
 ATT TGG TAC GAA AGC TAA TTC GTC GA

24

Lbc₁ with the sequence:

1320

TAA/TT AGG ATC TAC TGC ATT GCC GTA

1350 1380

AAG TGT AAT AAA TAA ATC TTG TTT CAA CTA AAA CTT GTT ATT AAA CAA GTT CCC TAT ATA

1410 1440

AAT GTT GTT TAA AAT AAG TAA ATT TCA TTG TAT TGG ATA AAC ACT TTT AAG TTA TAT ATT

1470 1500

5 TCC ATA TAT TTA CGT TTG TGA ATC ATA ATC GAT ACT TTA TTA AAA TAA ATT CCA AAT AAT

TTA TAC GTT TTA AAA ATT ATT TT

Lbc₂ with the sequence:

TAG/GAT CTA CTA TTG CCG TCA AGT
1140

GTA ATA AAT AAA TTT TGT TTC ACT AAA ACT TGT TAT TAA ACA AGT CCC CGA TAT ATA AAT
1170 1200

10 GTT GGT TAA AAT AAG TAA ATT ATA CGG TAT TGA TAA ACA ATC TTA AGT TTT ATA TAT AGT
1230 1260

TCC ATA TAC TAA AGT TTG TGA ATC ATA ATC GA
1290

and Lbc₃ with the sequence:

TAG/GAT CTA CAA TTG CCT TAA AGT GTA ATA AAT AAA
990 1020

TAT TAT TTC ACT AAA ACT TGT TAT TAA ACC AAG TTC TCG ATA TAA ATG TTG GTT AAA CTA
1050 1080

15 AGT AAA TTA TAT GGT ATT GGA TAA ACA ATC TTA AGC TT
1110

This sequence is positioned on the 0.9 Kb 3' flanking region used according to the invention. A particular embodiment of the invention is therefore

the use of sequences of this region exerting or mediating the regulation characterised by the invention of root nodule-specific promoter regions.

In a preferred embodiment of the method according to the invention a region is used of the coding sequence or intervening sequence of root nodule-specific genes, in particular sequences of the coding sequence or the intervening sequence capable of influencing the regulation of a promotor of the root nodule-specific genes or capable of influencing the yield of the desired gene product in another manner.

Examples of such coding sequences and intervening sequences are the four leghemoglobin genes of soybean, viz.

Lba with the sequence:

		120
		VAL
		ATG/GTT
	150	180
	ALA PHE THR GLU LYS GLN ASP ALA LEU VAL SER SER SER PHE GLU ALA PHE LYS ALA ASN	
	GCT TTC ACT GAG AAG CAA GAT GCT TTG GTG AGT AGC TCA TTC GAA GCA TTC AAG GCA AAC	
	210	240
	ILE PRO GLN TYR SER VAL VAL PHE TYR THR SER	
	ATT CCT CAA TAC AGC GTT GTG TTC TAC ACT TC/G TAA GTT TTC TCT CTA AGC ATG TGT CTT	
20	270	300
	CCA TTC TAT GTT TTT CTT TTG GAA ATT TGT TGT GTT TGA AAA AAG ATA TAT TGT TAA TGT	
	330	360
	GAG TGG TTT TGG TTT GAT TAA AAA TGA ATAG/G ATA CTG GAG AAA GCA CCT GCA GCA AAG GAC	
	390	420
	LEU PHE SER PHE LEU ALA ASN GLY VAL ASP PRO THR ASN PRO LYS LEU THR GLY HIS ALA	
	TTG TTC TCA TTT CTA GCA AAT GGA GTA GAC CCC ACT AAT CCT AAC CTC ACG GGC CAT GCT	
	450	480
	GLU LYS LEU PHE ALA LEU	
	GAA AAG CTT TTT GCA TTG/GTAA GTA TCA CCC AAC TAA AAT TAT AAC TAT TTT ATG TGA	
	510	540
	TTA ATT TTA AGA TTA AGC ATC ATG TAT TTT AAC ACT CTT AAA ACA TCA ATG AAC ATT AAT	
	570	600
25	TGT TTG AAT TGT ATT TTA TAT TTT TGC CAT ATC TTG AAC TAG GAA TAG TAT ATA AAT TTC	
	630	660
	TAT TAG TAT TTG TTG ATA ATT ATT TTT CTT TCA TAA CTA TCT TGT CAC ATA TTA TAT ATT	

```

                                690                                720
                                VAL ARG ASP SER ALA GLY GLN LEU LYS ALA SER GLY THR VAL VAL ALA
TTT TGA ATT GTAG/GTG CGT GAC TCA GCT GGT CAA CTT AAA GCA AGT GGA ACA GTG GTG GCT

                                750                                780
ASP ALA ALA LEU GLY SER VAL HIS ALA GLN LYS ALA VAL THR ASP PRO GLN PHE VAL
GAT GCC GCA CTT GGT TCT GTT CAT GCC CAA AAA GCA GTC ACT GAT CCT CAG TTC GTG/GT

                                810                                840
ATG ATA AAT AAT GAA ATG TTA TAA TAA ATT ATG CAT ACT TCA ATT TTT CAT GGA GCA GTA

                                870                                900
TAA TGA TCA ACA CAC ACT TCT TTT GTT TCA TGC ATT TGA TAA CTA CAA TCT TAA AAT GTT

                                930                                960
5 GCA ATC TTA AAA ATA GTA TTA AAA ATA TAA CAT TTA ATT AGC TCA TCA ATA TTT TTC TGT

                                990                                1020
TGC AAT TTT TTA TGA AAA AAT TAT AAT TAT GAA TTC TTT GAG CAA TGT TTA ATT AAA AAA

                                1050                                1080
TTG ATT TAA TAA TGA AAT AAC TAA GCT ACC TCT GTC TCG TTT TTC ATT TAA ACT ATG ACA

                                1110                                1140
TAA ACA ATG AAT AAA GTA AAC TAA ACC ATG ACA TGT TTA TTT TTG AAT GAG GTT ATT AAT

                                1170                                1200
AAT TTT TTT TCA CTA TCT ATT GCA ATG TTC ATT GAT TAT CAA TTA TCT TGG TTG CAT TGA

                                1230                                1260
10 TTC TCT CGA TTT TTT TCT TGA GGT TAA GCT TCA GTT CAA TAT ATA TTC ATT TTT TGA TAA

                                1290                                1320
AAA AAA ATA GTA CAA TAT ATT TTC AAT TAG CTG ATC ATA TTT ATT TAA GTT CAA CTT AAA

                                1350                                1380
ATT TTA TAG ATG TTA ATT GAT ATA ATT TGT TGA GAT GAT GAG AAG ACC AAT ACC ATT ACG

                                1410                                1440
TAC TCT TTT GAA AGT GTT ATA TGG AAT TTA ATT ATA AGG AAA AAT GTA AGA GCT AAA CCA

                                1470                                1500
                                VAL VAL LYS GLU ALA LEU LEU LYS THR ILE LYS ALA ALA VAL
15 TTG CTG ATG ATT TTG AAG/GTG GTT AAA GAA GCA CTG CTG AAA ACA ATA AAG GCA GCA GTT

                                1530                                1560
GLY ASP LYS TRP SER ASP GLU LEU SER ARG ALA TRP GLU VAL ALA TYR ASP GLU LEU ALA
GGG GAC AAA TGG AGT GAC GAG TTG AGC CGT GCT TGG GAA GTA GCC TAC GAT GAA TTG GCA

ALA ALA ILE LYS LYS ALA
GCA GCT ATT AAG AAG GCA TAA

```

The amino acid sequence of the Lba protein is indicated above the coding sequence,

Lbc₁ with the sequence:

0249676

27

180
GLY
ATG/GCT

210 240
 ALA PHE THR GLU LYS GLN GLU ALA LEU VAL SER SER SER PHE GLU ALA PHE LYS ALA ASN
 GCT TTC ACT GAG AAG CAA GAG GCT TTG GTG AGT AGC TCA TTC GAA GCA TTC AAG GCA AAC

270 300
 ILE PRO GLN TYR SER VAL VAL PHE TYR ASN SER
 ATT CCT CAA TAC AGC GTT GTG TTC TAC AAT TC/GTAA GTT TTC TCT ATA AGC ATG TGT CTT

330 360
 TCA TTC TAT GTT TTT CTT CTG GAA ATT TTT TGT GTT TGA AAA AAG ATA TAT ATA TAT ATA

390 420
 5 TAT ATA TAT ATA TAT ATA TAT ATA TAT ATA TAT TTT GTT AAT GTG AGT GGT TTT

450 480
 ILE LEU GLU LYS ALA PRO ALA ALA LYS ASP LEU PHE SER
 GGT TTG ATT AAA AAT AAA TAG/GATT CTG GAG AAA GCA CCT GCA GCA AAG GAC TTG TTC TCA

510 540
 PHE LEU ALA ASN GLY VAL ASP PRO THR ASN PRO LYS LEU THR GLY HIS ALA GLU LYS LEU
 TTT CTA GCA AAT GGA GTA GAC CCC ACT AAT CCT AAG CTC ACG GGC CAT GCT GAA AAG CTT

570 600
 PHE ALA LEU
 TTT GCA TTG/GT AAG TAT CAG CCA ACT AAA ATT ATA ACT ATT TTA TGT GAT TAA TTT TAA

630 660
 GAT TAA ACA TCA TGT ATT TTA ACA CTC TTA AAA TAT CAA TGA ACA TTA ATT TTT TGA ATT

690 720
 10 GTA TTT TAT ATT TTT ACC ATA TCT TGA ACT AGG AAT AAT ATA TAA ATT TCT ATT AGT ATT

750 780
 TGT TGG TAA TTA CAT ATA TAT ATA TAT ATA TAA TCC TTG TGA TAA TTA TTT TTC GAA TTT

810 840
 VAL ARG ASP SER ALA GLY GLN LEU LYS THR ASN GLY THR VAL VAL ALA ASP ALA ALA
 GTAG/GTG CGT GAC TCA GCT GGT CAA CTT AAA ACA AAT GGA ACA GTG GTG GCT GAT GCT GCA

870 900
 LEU VAL SER ILE HIS ALA GLN LYS ALA VAL THR ASP PRO GLN PHE VAL
 CTT GTT TCT ATC CAT GCC CAA AAA GCA GTC ACT GAT CCT CAG TTC GTG/GT ATG ATA AAT

930 960
 AAT ACT AGT AAA ATG TTA CAA TAA ATG CAA ACT TAA GTT TTA CGT ACA TAG TGA TCA TGA

990 1020
 15 CTT CAT GCA TGG CTA TTA TTT TTT CAT ATT TAT TGA AGT CAA CTT AAA ATT TTG TAA ATA

1050 1080
 CAG ATC GAT GCT AGT AAT TTG TTG AGA TCA TGA GAA AAC GTA CCA CTA CTC CAA TAG CAT

1110 1140
 TAC TCA TTT TGA AAA TTG TAT AAC TGT GAT CTA ATT ATA AGG AAA AAG TGT ATA TAA GAG

1170 1200
 CTA ATC CAT TAT TAA TGT TTT TTA TAT TTT GTAG/GTG GTT AAA GAA GCA CTG CTG AAA ACA
 VAL VAL LYS GLU ALA LEU LEU LYS THR

1230 1260
 ILE LYS GLU ALA VAL GLY GLY ASN TRP SER ASP GLU LEU SER SER ALA TRP GLU VAL ALA
 ATA AAG GAA GCT GTT GGC GGC AAT TGG AGT GAC GAA TTG AGC AGT GCT TGG GAA GTA GCC

1290
 20 TYR ASP GLU LEU ALA ALA ALA ILE LYS LYS ALA
 TAT GAT GAA TTG GCA GCA GCA ATT AAA AAG GCA TAA

The amino acid sequence of the Lbc₁ protein is indicated above the coding sequence,

0249676

28

Lbc₂ with the sequence:

GLY
G/GGT
180

ALA PHE THR GLU LYS GLN GLU ALA LEU VAL SER SER SER PHE GLU ALA PHE LYS ALA ASN
GCT TTC ACT GAG AAG CAA GAG GCT TTG GTG AGT AGC TCA TTC GAA GCA TTC AAG GCA AAC
210 240

ILE PRO GLN TYR SER VAL VAL PHE TYR THR SER
ATT CCT CAA TAC AGC GTT GTG TTC TAC ACT TC/GTA AGT TTT CTC TTA AAG CAT GTA TCT
270 300

5 TTC ATT CTC TGT TTT TCC TTT CGA CAT TTT TTG TGT TTG AAA AGA GAT AGT GTC AAT GTG
330 360

AGT GGG TAT TTT TTT TTA TTA AAA ATT AAC AG/G ATA CTG GAG AAA GCA CCC GCA GCA AAG
390 420

ASP LEU PHE SER PHE LEU SER ASN GLY VAL ASP PRO SER ASN PRO LYS LEU THR GLY HIS
GAC TTG TTC TCG TTT CTA TCT AAT GGA GTA GAT CCT AGT AAT CCT AAG CTC ACG GGC CAT
450 480

ALA GLU LYS LEU PHE GLY LEU
GCT GAA AAG CTT TTT GGA TTG/GTA AGT ATC ATC CAA CTA AAA TTA TAG CTA TTT TAT GTG
510 540

10 ATT AAT TTT AAG ATT AAA CAT GTA TTT AAC ACT CTT AAA CAT GTA TTT AAC ACT CTT AAG
570 600

ATT AAA CAT GTA TTT AAC TAA AAC ATG TAT TTG CTG ATT ATT TTT TTT TTA TAA TTA TCT
630 660

TGT CAC ATA TTA TAT ATT TTT TGA ATT GTA VAL ARG ASP SER ALA GLY GLN LEU LYS ALA
690 720

ASN GLY THR VAL VAL ALA ASP ALA ALA LEU GLY SER ILE HIS ALA GLN LYS ALA ILE THR
AAT GGA ACA GTA GTG GCT GAT GCC GCA CTT GGT TCT ATC CAT GCC CAA AAA GCA ATC ACT
750 780

15 ASP PRO GLN PHE VAL
GAT CCT CAG TTC GTG/GT ATG ATA AAT AAT AAA ATG TTA CAA TAA ATG CAC ATA TAC TTA
810 840

AAT TTT ACA TGG TGC AGT GTT ATG ATC ATC ATT TTT GTT TAG TAA TGA ATT TAC TTA AAA
870 900

TCT TAA ATT ATG TAC TTT TTG AAA GTT TTA TAT GGA ATT TTA ATT ATA GGG AAA AAT GTA
930 960

AGA GCT AAT CCA TTA GTG ATG TTT TGT CTG VAL VAL LYS GLU ALA LEU LEU LYS THR
990 1020

LE LYS GLU ALA VAL GLY ASP LYS TRP SER ASP GLU LEU SER SER ALA TRP GLU VAL ALA
ATA AAG GAG GCA GTT GGG GAC AAA TGG AGT GAT GAA TTG AGC AGT GCT TGG GAA GTA GCC
1050 1080

20 TYR ASP GLU LEU ALA ALA ALA ILE LYS LYS ALA PHE
TAT GAT GAA TTG GCA GCA GCT ATT AAG AAG GCA TTT TAC
1110

0249676

29

The amino acid sequence of the Lbc₂ protein is indicated above the coding sequence,

and Lbc₃ with the sequence:

```

                                GLY ALA PHE THR ASP
                                G/GGT GCT TTC ACT GAT
                                120
LYS GLN GLU ALA LEU VAL SER SER SER PHE GLU ALA PHE LYS THR ASN ILE PRO GLN TYR
5 AAG CAA GAG GCT TTG GTG AGT AGC TCA TTT GAA GCA TTC AAG ACA AAC ATT CCT CAA TAC
  150 180
SER VAL VAL PHE TYR THR SER
AGT GTT GTG TTC TAC ACC TC/GTA AGT ATT CTA TCT AAA TTA TGT GTC TTA TTG TAT GTT
  210 240
TAA CTT TCG TGG TTT GTT GTG TTT GAA AAA AAG ATA TAT ATT GTT AAT GTG AGT GGT TTT
  270 300
                                ILE LEU GLU LYS ALA PRO VAL ALA LYS ASP LEU PHE SER
GGT TTG ACT AAA AAT GAA TAG/G ATA CTG GAG AAA GCA CCT GTA GCA AAG GAC TTG TTC TCA
  330 360
10 PHE LEU ALA ASN GLY VAL ASP PRO THR ASN PRO LYS LEU THR GLY HIS ALA GLU LYS LEU
   TTT CTA GCT AAT GGA GTA GAC CCC ACT AAT CCT AAG CTC ACG GGC CAT GCT GAA AAA CTT
     390 420
PHE GLY LEU
TTT GGA TTG/GT AAG TAT CCA GCC TAC TAA AAT TAA AAT CCT ATT AGT ATT TTT TAT TAT
  450 480
                                VAL ARG ASP SER
TTT TCT TCC ATG ATT GTC TTG TCA CAT ATT ATA TAT TTT TTG AAT TAT AG/GTA CGT GAT TCA
  510 540
ALA GLY GLN LEU LYS ALA SER GLY THR VAL VAL ILE ASP ALA ALA LEU GLY SER ILE HIS
GCT GGT CAA CTT AAA GCA AGT GGA ACA GTG GTG ATT GAT GCC GCA CTT GGT TCT ATC CAT
  570 600
15 ALA GLN LYS ALA ILE THR ASP PRO GLN PHE VAL
   GCC CAA AAA GCA ATC ACT GAT CCT CAA TTT GTG/G TAT GAT AAA TAA TGA AAA GCT ACA
     630 660
ATA AAT GCA CAA ATA CTT AAT TTT ACA TAG TGC AGT GCT ATA TGA TCA TCA CTT TTG CTT
  690 720
AGT AAT GAA TTT ACT TTT TTT TTT TAC AGA AGT AAT GGA TTT ACT TAA AAT CTT AAA TTA
  750 780
TGT ACT TCT TTA AAG AGT TTT GTA TGG AAT TTT AAT TAT AGG AAA AAT GTA AGA GCT AAA
  810 840
                                VAL VAL LYS GLU ALA LEU LEU LYS THR ILE LYS GLU ALA
CCA TTG CTG ATG ATT TCG AAG/GTG GTT AAA GAA GCA CTG CTG AAA ACA ATA AAG GAG GCA
  870 900
20 VAL GLY ASP LYS TRP SER ASP GLU LEU SER SER ALA TRP GLU VAL ALA TYR ASP GLU LEU
   GTT GGG GAC AAA TGG AGT GAC GAG TTG AGC AGT GCT TGG GAA GTA GCC TAT GAT GAA TTG
     930 960
ALA ALA ALA ILE LYS LYS ALA PHE
GCA GCA GCT ATT AAG AAG GCA TTT TAG

```

The amino acid sequence of the Lbc₃ protein is indicated above the coding sequence.

The present invention furthermore deals with a novel DNA fragment comprising an inducible plant promoter to be used when carrying out the method according to the invention, said DNA fragment being characterised by being identical with, derived from or comprising a 5' flanking region of root nodule-specific genes. Examples of such DNA fragments are DNA fragments being identical with, derived from or comprising a 5' flanking region of plant leghemoglobin genes. Preferred examples are according to the invention DNA fragments being identical with, derived from or comprising a 5' flanking region of the four soybean leghemoglobin genes, viz.:

Lba with the sequence:

	GAGATACATT	ATAATAATCT	CTCTAGTGTC	TATTTATTAT	TTTATCTGGT
	GATATATACC	TTCTCGTATA	CTGTTATTTT	TTCAATCTTG	TAGATTTACT
20	TCTTTTATTT	TTATAAAAAA	GACTTTATTT	TTTTAAAAAA	AATAAAGTGA
	ATTTTGAAAA	CATGCTCTTT	GACAATTTTC	TGTTTCCTTT	TTCATCATTG
	GGTTAAATCT	CATAGTGCCT	CTATTCAATA	ATTGCGGCTC	AATTTAATTA
	GTAGAGTCTA	CATAAAATTT	ACCTTAATAG	TAGAGAATAG	AGAGTCTTGG
	AAAGTTGGTT	TTTCTCGAGG	AAGAAAGGAA	ATGTTAAAAA	CTGTGATATT
	TTTTTTTTTG	ATTAATAGTT	ATGTTTATAT	GAAAACTGAA	AATAAATAAA
25	CTAACCATAT	TAAATTTAGA	ACAACACTTC	AATTATTTTT	TTAATTTGAT
	TAATTAAAAA	ATTATTTGAT	TAAATTTTTT	AAAAGATCGT	TGTTTCTTCT
	TCATCATGCT	GATTGACACC	CTCCACAAGC	CAAGAGAAAC	ACATAAGCTT
	TGGTTTTCTC	ACTCTCCAAG	CCCTCTATAT	AAACAAATAT	TGGAGTGAAG
	TTGTTGCATA	ACTTGCATCG	AACAATTAAT	AGAAATAACA	GAAAATTAAA
	AAAGAAATAT	G,			

Lbc₁ with the sequence

```

TTCTCTTAAT ACAATGGAGT TTTTGTGAA CATACATACA TTTAAAAAAA
AATCTCTAGT GTCTATTTAC CCGGTGAGAA GCCTTCTCGT GTTTTACACA
CTTTAATATT ATTATATCCT CAACCCACACA AAAAAGAATA CTGTTATATC
5 TTTCCAAACC TGTAGATTTA TTTATTTATT TATTTATTTT TACAAAGGAG
ACTTCAGAAA AGTAATTACA TAAAGATAGT GAACATCATT TTATTTATTA
TAATAAACTT TAAAATCAAA CTTTTTTATA TTTTTTGTTA CCCTTTTCAT
TATTGGGTGA AATCTCATAG TGAAGCCATT AAATAATTTG GGCTCAAGTT
TTATTAGTAA AGTCTGCATG AAATTTAACT TAACAATAGA GAGAGTTTTC
GAAAGGGAGC GAATGTTAAA AAGTGTGATA TTATATTTTA TTTGATTAA
TAATTATGTT TACATGAAAA CATACAAAAA AATACTTTTA AATTCAGAAT
AATACTTAAA ATATTTATTT GCTTAATTGA TTAAGTAAA ATTATTTGAT
10 TAGGATTTTG AAAAGATCAT TGGCTCTTCG TCATGCCGAT TGACACCCTC
CACAGCCAA GAGAACTTA AGTTGTAAAC TTTCTCACTC CAAGCCTTCT
ATATAAACAT GTATTGGATG TGAAGTTATT GCATAACTTG CATTGAACAA
TAGAAAAATA CAAAAAAAG TAAAAAAGTA GAAAAGAAAT ATG,

```

Lbc₂ with the sequence:

```

TCGAGTTTTT ACTGAACATA CATTTATTAA AAAAAACTCT CTAGTGTCCA
TTTATTCGGC GAGAAGCCTT CTCGTGCTTT ACACACTTTA ATATTATTAT
15 ATCCCCACCC CCACCAAAAA AAAAAAAACT GTTATATCTT TCCAGTACAT
TTATTTCTTA TTTTACAAA GGAACTTCA CGAAAGTAAT TACAAAAAAG
ATAGTGAACA TCATTTTTTT AGTTAAGATG AATTTTAAAA TCACACTTTT
TTATATTTTT TTGTTACCCT TTTCATTATT GGGTGAAATC TCATAGTGAA
ACTATTAAAT AGTTTGGGCT CAAGTTTAT TAGTAAAGTC TGCATGAAAT
TTAACTTAAT AATAGAGAGA GTTTTGGAAG GGTAACGAAT GTTAGAAAGT
GTGATATTAT TATAGTTTAA TTTAGATTAA TAATTATGTT TACATGAAAA
TTGACAATTT ATTTTAAAA TTCAGAGTAA TACTTAAATT ACTTATTTAC
20 TTTAAGATTT TGAAAAGATC ATTTGGCTCT TCATCATGCC GATTGACACC
CTCCACAAGC CAAGAGAAAC TTAAGTTGTA ATTTTCTAA CTCCAGCCT
TCTATATAAA CACGTATTGG ATGTGAAGTT GTTGATAAC TTGCATTGAA
CAATAGAAAT AACACAAAG AAAATAAGTG AAAAAAGAAA TATG,

```

and Lbc₃ with the sequence:


```

TATGAAGATT AAAAAATACA CTCATATATA TGCCATAAGA ACCAACA AAAA
GTACTATTTA AGAAAAGAAA AAAAAAACCT GCTACATAAT TTCCAATCCT
GTAGATTTAT TTCTTTTATT TTTATAAAGG AGAGTTAAAA AAATTACAAA
ATAAAAATAG TGAACATCGT CTAAGCATTT TTATATAAGA TGAATTTTAA
AAATATAATT TTTTGTCTA AATCGTATGT ATCTTGTCTT AGAGCCATTT
5 TTGTTTAAAT TGGATAAGAT CACACTATAA AGTTCTTCCT CCGAGTTTGA
TATAAAAAAA ATTGTTTCCC TTTTGATTAT TGGATAAAAT CTCGTAGTGA
CATTATATTA AAAAAATTAG GGCTCAATTT TTATTAGTAT AGTTTGCATA
AATTTTAACT TAAAAATAGA GAAAATCTGG AAAAGGGACT GTTAAAAAGT
GTGATATTAG AAATTTGTCG GATATATTAA TATTTTATTT TATATGGAAA
CTAAAAAAT ATATATTAAA ATTTTAAATT CAGAATAATA CTTAAATTAT
TTATTTACTG AAAATGAGTT GATTTAAGTT TTTGAAAAGA TGATTGTCTC
TTCACCATAC CAATTGATCA CCCTCCTCCA ACAAGCCAAG AGAGACATAA
10 GTTTTATTAG TTATTCTGAT CACTCTTCAA GCCTTCTATA TAAATAAGTA
TTGGATGTGA AGTTGTTGCA TAACTTGCAT TGAACAATTA ATAGAAATAA
CAGAAAAGTA GAAAAGAAAT ATG.

```

Another example of a preferred DNA fragment according to the invention is a DNA fragment which is 15 identical with, derived from or comprises 5' flanking regions of the Lbc3-5'-3'CAT gene with the sequence

```

TATGAAGATT AAAAAATACA CTCATATATA TGCCATAAGA ACCAACA AAAA
GTACTATTTA AGAAAAGAAA AAAAAAACCT GCTACATAAT TTCCAATCCT
GTAGATTTAT TTCTTTTATT TTTATAAAGG AGAGTTAAAA AAATTACAAA
20 ATAAAAATAG TGAACATCGT CTAAGCATTT TTATATAAGA TGAATTTTAA
AAATATAATT TTTTGTCTA AATCGTATGT ATCTTGTCTT AGAGCCATTT
TTGTTTAAAT TGGATAAGAT CACACTATAA AGTTCTTCCT CCGAGTTTGA
TATAAAAAAA ATTGTTTCCC TTTTGATTAT TGGATAAAAT CTCGTAGTGA
CATTATATTA AAAAAATTAG GGCTCAATTT TTATTAGTAT AGTTTGCATA
AATTTTAACT TAAAAATAGA GAAAATCTGG AAAAGGGACT GTTAAAAAGT
GTGATATTAG AAATTTGTCG GATATATTAA TATTTTATTT TATATGGAAA
25 CTAAAAAAT ATATATTAAA ATTTTAAATT CAGAATAATA CTTAAATTAT
TTATTTACTG AAAATGAGTT GATTTAAGTT TTTGAAAAGA TGATTGTCTC
TTCACCATAC CAATTGATCA CCCTCCTCCA ACAAGCCAAG AGAGACATAA
GTTTATTAG TTATTCTGAT CACTCTTCAA GCCTTCTATA TAAATAAGTA
TTGGATGTGA AGTTGTTGCA TAACTTGCAT TGAACAATTA ATAGAAATAA
CAGAAAAGTA GAATTCCTAAA ATG

```

30 Still another example of such a DNA fragment ac-

cording to the invention is a DNA fragment which is identical with, derived from or comprises 5' flanking regions of the N23 gene with the sequence

```

      10      20      30      40      50      60      70
GAATTCGAGCTCGCCCGGGGATCGATCCTCTAGAGTCGACCTGCAGCCCAAGCTTGGATCAATCAATTAA
  ECORI                               SalI
      80      90     100     110     120     130     140
5  TTCTATTGAGACACGATTGAACAATTTTACATTATGAGACTATTTTGGTTTTTATTTGATCCAAA

      150     160     170     180     190     200     210
AAATTAAAGCTTTAGATGATGATGAATTGAANNAATATTGTATTAAATNNTGAAAAGTTNNNNNGGTTTA

      220     230     240     250     260     270     280
ATGAATGCTATGATATTGATGGTCTTGATNTATTNNCAGAATTGAAAGTATTAAGAGAAGTGTTAAGAAA

      290     300     310     320     330     340     350
10 AGAAGTTAGCACACCAATAGAAGTATTGAGTTATATTAACCTTTAGATTCTTTCAAATGTTTACATTG

      360     370     380     390     400     410     420
CATATAGAATTTTATTGACAATCCTTATAACAGTTGCTACTGTTGAAAGACGTTCTTCAAATTAATAATT

      430     440     450     460     470     480     490
ACTTAAATCATATCTAAATCAACAATGTTACAAGATAGATTGAATGAGTTAGTTATTTTATCTATTGAA

      500     510     520     530     540     550     560
15 AGTAAAGTGTTAGAATTGTTTGATTATAAACTCTGATAAATGATTTGTCAGTTAAAAAACTAGAAGAT

      570     580     590     600     610     620     630
TAATATAAAATTGATATTTTATATAATATATTAAGTCTCTTAAATTTCTTGTAAGAAAAAGACATTTT

      640     650     660     670     680     690     700
AAATAATAAAATAAGCAACTCTTAATTTTAAAGAACATCCCTTTGTTAAACCGAATCTTCCATAATGT

      710     720     730     740     750     760     770
20 AAAAATTAATGCTTGATGGAAGTTTTTAATTTGTTCTACTCAATACTCAAAGGGTTGTAATATTTTTT

      780     790     800     810     820     830     840
TATCATTTATATGTTGTAAATATGAATGCACTAGTAATTAGTTTAAATGATAAAATATATTCTACAGATAT

      850     860     870     880     890     900     910
ATTTCTGTCTCTTGGCAACTCGTGAGAATTGAATATATTATAAAGATGAAAGGTCGTTACAATTTTTTT

      920     930     940     950     960     970     980
25 AGAATAAATATTTATATACAATTCCTAGATTTTGTATAAAATTCACATATTGTATGAGTATAAATACAT

      990    1000    1010    1020    1030    1040    1050
GAGCACACACCAAACCTAGTCTCAAATTAAGTAAGGTGCTAATTATTAGCGGCTAGCTAAGTAACCAAGTA
                               DdeI

ATTAATG

```

The invention relates furthermore to any plasmid to be used when carrying out the method according to the invention and characterised by comprising a DNA fragment containing an inducible plant promoter
5 as herein defined. Particular examples of suitable plasmids according to the invention are pAR11, pAR29, pAR30, and N23-CAT, cf. Examples 3, 4, and 11. These plasmids allow recombination into the A. rhizogenes T DNA region.

- 10 The invention relates furthermore to any Agrobacterium strain to be used in connection with the invention and characterised by comprising a DNA fragment comprising an inducible plant promoter of root nodule-specific genes built into the T DNA
15 region and therefore capable of transforming the inducible promoter into plants. Particular examples of bacterium strains according to the invention are the A. rhizogenes strains AR1127 carrying pAR29, AR1134 carrying pAR30, AR1000 carrying pAR11, and
20 AR204-N23-CAT carrying N23-CAT.

It is obvious that the patent protection of the present invention is not limited by the embodiments stated above.

Thus the invention employs not exclusively 5' flanking regions of soybean leghemoglobin genes. It is
25 well-known that the leghemoglobin genes of all leguminous plants have the same function, cf. Appleby (1974) in The Biology of Nitrogen Fixation, Quispel. A. Ed. North-Holland Publishing Company, Amsterdam, Oxford, pages 499-554, and concerning the
30 kidney bean PvLb1 gene it has furthermore been

proved that a high degree of homology exists with the sequences of the soybean Lbc₃ gene. It is also known that the expression of other root nodule-specific genes is regulated in a similar manner
5 like the leghemoglobin genes. The invention includes thus the use of 5' flanking regions of leghemoglobin genes or other root nodule-specific genes of all plants in case the use of such DNA fragments makes the expression of a desired gene product the subject
10 matter of the regulation characterised by the present invention.

The present invention allows also the use of such fragments of any origin which under natural conditions exert or mediate the regulation characterised by the present invention. The latter applies
15 especially to such fragments which can be isolated from DNA fragments from gene libraries or genomes through hybridization with labelled sequences of 5' flanking regions of soybean leghemoglobin genes.

20 It is well-known that it is possible to alter nucleotide sequences of non-important sub-regions of 5' flanking regions without causing an alteration of the promoter activity and the regulation. It is also well-known that an alteration of sequences of
25 important subregions of 5' flanking regions renders it possible to alter the binding affinities between nucleotide sequences and the factors or effector substances necessary or responsible for the transcription initiation and the translation initiation
30 and consequently to improve the promoter activity and/or the regulation. The present invention includes, of course, also the use of DNA fragments

containing such altered sequences of 5' flanking regions, and in particular DNA fragments can be mentioned which have been produced by recombining sequences of 5' flanking regions of any gene with
5 5' flanking regions of root nodule-specific genes provided the use of such DNA fragments subjects the expression of a desired gene product to the regulation characterised by the present invention.

It should be noted that the transformation of micro-
10 organisms is carried out in a manner known per se, cf. e.g. Maniatis et al., (1982), Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Laboratory.

The transformation of plant cells, i.e. introduction of plasmid DNA into plant cells, is also carried
15 out in a manner known per se, cf. Sambrook et al., (1983), EMBO J. 2, 2143-2150.

Cleavage with restriction endonucleases and digestion with other DNA modifying enzymes are well-known techniques and are carried out as recommended
20 by the suppliers.

The Agrobacterium rhizogenes 15834 rif^R was used as a typical representative of A. rhizogenes: see White et al., I.Bact., Vol. 141 (1980), 1134-1141.

Example 1

25 Sequence determination of 5' flanking regions of soybean leghemoglobin genes

From a soybean gene library the four soybean leg-

hemoglobin genes Lba, Lbc₁, Lbc₂, and Lbc₃ are provided as described by Jensen, E.Ø. et al., Nature Vol. 291, No. 3817, 677-679 (1981). The genetically stable in-bred invariable soybean species "Glycine max.var.Evans" was used as a starting material for the isolation of the DNA used for the construction of said gene library. The 5' flanking regions of the four soybean leghemoglobin genes are isolated, as described by Jensen, E.Ø., Ph D Thesis, Institut for Molekylær Biologi, Århus Universitet (1985), and the DNA sequences determined by the use of the dideoxy method as described by Sanger, F., J. Mol. Bio. 143, 161-178 (1980) and indicated in the sequence scheme.

15 Example 2

Construction of Lbc₃-5'-3'-CAT

The construction has been carried out in a sequence of process steps as described below:

a) Sub-cloning the Lbc₃ gene

20 The Lbc₃ gene was isolated on a 12Kb EcoRI restriction fragment from a soybean DNA library, which has been described by Wiborg et al., in Nucl. Acids Res. (1982) 10, 3487. A section of the fragment is shown at the top of the attached Scheme 2. This
25 fragment was digested by the enzymes stated and then ligated to pBR322 as indicated at the Scheme. The resulting plasmids Lbc₃HH and Lbc₃HX were subsequently digested by PvuII and religated, which result d in two plasmids called pLpHH and pLpHX.

b) Sub-cloning 5' flanking sequences from the Lbc₃ gene

For this purpose pLpHH was used as shown in the attached Scheme 3. This plasmid was opened by means of PvuII and treated with exonuclease Bal31. The reaction was stopped at various times and the shortened plasmids were ligated into fragments from pBR322. These fragments had been treated in advance as shown in Scheme 3, in such a manner that in one end they had a DNA sequence TTC ---
AAG ---

After the ligation a digestion with EcoRI took place, and the fragments containing 5' flanking sequences were ligated into EcoRI digested pBR322. 15 These plasmids were transformed into E. coli K803, and the plasmids in the transformants were tested by sequence analysis. A plasmid, p213 5'Lb, isolated from one of the transformants, contained a 5' flanking sequence terminating 7 bp before the Lb ATG 20 start codon in such a manner that the sequence is as follows:

2Kb
-5' flanking --- AAAGTAGAATTC
Lbc₃ sequence

25 E.coli K803 is a typical representative of the E. coli K12 recipient strains.

c) Sub-cloning 3' flanking region of the Lbc₃ gene

For this purpose pLpHX was used which was digested by XhoII. The ends were partially filled out and excess single-stranded DNA was removed with S1 nuclease, as shown in the attached Scheme 4. The 5 fragment shown was ligated into pBR322 which had been pretreated as shown in the Scheme. The construction was transformed into E. coli K803. One of the transformants contained a plasmid called Xho2a-3'Lb. As the XhoII recognition sequence is 10 positioned immediately after the Lb stop codon, cf. Scheme 2, the plasmid contained about 900 bp of the 3' flanking region, and the sequence started with GAATTCTACAA---.

The construction of Lb promoter cassette

15 An EcoRI/SphI fragment from Xho2a-3'Lb was mixed with a BamHI/EcoRI fragment from p213-5'Lb. These two fragments were ligated via the BamHI/SphI cleavage sites into a pBR322 derivative where the EcoRI recognition sequence had been removed, cf. Scheme 20 4. The ligated plasmids were transformed into E. coli K803. A plasmid in one of the transformants contained the correct fragments, and it was called pEJLb 5'-3'-1.

Construction of the Lbc3 5'3'-CAT gene

25 The CAT gene of pBR322 was isolated on several smaller restriction fragments, as shown in the attached Scheme 5. The 5' coding region was isolated as an AluI fragment which was subsequently ligated into pBR322, treated as stated in the Scheme. This

was transformed into E. coli K803. Several transformants contained the correct plasmid. One was taken out and called Alu11. The 3' coding region was isolated on a TaqI fragment. This fragment was
5 treated with exonuclease Bal31, whereafter EcoRI linkers were added. Then followed a digestion with EcoRI and a ligation to EcoRI digested pBR322. The latter was transformed into E. coli K803 and the transformants were analysed. A plasmid, Taq 12,
10 contained the 3' coding region of the CAT gene plus 23 bp 3' flanking sequences subsequently terminating in the following sequence CCGGAATTC. Subsequently the following fragments were ligated together to EcoRI digested
15 pEJLb5'-3'-1: EcoRI/PvuII fragment from AluI, PvuII/DdeI fragment from pBR322 and DdeI/EcoRI fragment from Taq 12. This ligation mixture was transformed into E. coli K803. Several transformants contained the correct plasmid. One was taken out
20 and was called pEJLb 5'-3' CAT 15.

Example 3

a.

Cloning and integration of the soybean Lbc₃-5'-3'-CAT gene.

25 Two EcoRI fragments (No. 36 and No. 40) of the T_L-DNA region of A. rhizogenes 15834 pRi plasmid was used as "integration sites". Thus the Lbc₃-5'-3'-CAT gene was subcloned (as 3,6 Kb BamHI/SalI fragment) into two vectors pAR1 and pAR22 carrying the
30 above EcoRI fragments. The resulting plasmids pAR29

and pAR30 were separately mobilized into A. rhizogenes 15834 rif^R using a plasmid helper system; see E. van Haute et al. (1983), EMBO J. 3, 411-417. Neither pAR29 nor pAR30 can replicate in Agrobacterium. Therefore the selection by means of rifampicin 100 µg/ml and the plasmid markers spectinomycin 100 µg/ml, streptomycin 100 µg/ml or kanamycin 300 µg/ml will select A. rhizogenes bacteria having integrated the plasmids via homologous recombination through the EcoRI fragments 36 or 40. The structure of the resulting T_L-DNA regions - transferred to the transformed plant lines L5-9 and L6-23 - has been indicated at the bottom of the attached Scheme 6. In this Scheme is furthermore for the L6-23 line shown the EcoRI and HindIII fragments carrying the Lbc₃-5'-3'-CAT gene and therefore hybridizing to radioactively labelled Lbc₃-5'-3'-CAT DNA used as a probe, cf. Example 4a.

20b.

Cloning and integration of the soybean Lbc₃ gene.

The EcoRI fragment No. 40 has here been used as "integration site". The Lbc₃ gene was therefore sub-cloned (as a 3,6 Kb BamHI fragment into the 25pAR1 vector and transferred into the T_L-DNA region as stated in a. The structure of the T_L-DNA region, transferred to the transformed plant line L8-35, has been shown at the bottom of the attached Scheme 7. This Scheme furthermore shows the EcoRI and HindIII fragments carrying the Lbc₃ gene and there-


0249676

42

fore hybridizing with radioactively labelled Lbc₃
DNA used as a probe, cf. Example 4b.

Example 4.a.


Demonstration of the soybean Lbc₃-5'-3'-CAT gene in transformed plants of bird's-foot trefoil.



DNA extracted from transformed lines (L6-'23) or untransformed control plants and cleaved by the restriction enzymes EcoRI and HindIII was analyzed by Southern-hybridization. Radioactively labelled Lbc₃-5'-3'-CAT gene was used as a probe for demonstrating corresponding sequences in the transformed lines. The bands marked with numbers correspond to restriction fragments constituting parts of the Lbc₃-5'-3'-CAT gene as stated in the restriction map (Scheme 6) of Example 3a.

b.


Demonstration of the soybean Lbc₃ gene of transformed plants of bird's-foot trefoil.



DNA extracted from transformed lines (L8-35) or untransformed control plants and cleaved by the restriction enzymes EcoRI and HindIII was analyzed by Southern-hybridization. Radioactive Lbc₃ gene 5 was used as a probe for detecting corresponding sequences in the transformed lines. The bands marked with numbers correspond to restriction fragments constituting parts of the Lbc₃ gene as stated in the restriction map (Scheme 7) of Example 3b.

Example 5a.

Expression of the Lbc₃-5'-3'-CAT gene in various tissues of bird's-foot trefoil.



The activity of the chloroamphenicol acetyl transferase (CAT) enzyme is measured as the amount of acetylated chloroamphenicol (AcCm) produced from ^{14}C -chloroamphenicol. In (a) the acetylated forms 1AcCm and 3AcCm appear, which have been separated from Cm through thin-layer chromatography in chloroform/methanol (95:5). The columns 1-3 show that no CAT activity occurs in root (R), nodule (N), as well as leaves + stem (LS) of untransformed plants of bird's-foot trefoil. The columns 4-6 and 7-9 show the CAT activity in corresponding tissues of Lbc₃-5'-3'-CAT transformed L6-23 and L5-9 plants. The conversion of chloroamphenicol in columns 5 and 8 shows the organ-specific expression of the Lbc₃-5'-3'-CAT gene in root nodules. The columns 10-12 show the lack of CAT activity in plants transformed with the Lbc₃ gene.

b.

Table


20	L6-23		L5-9	
	CAT activity		CAT activity	
Root	0		0	
Nodule	68830 cpm/ μg protein.h		154,000 cpm/ μg protein.h	
Leaves +				
25 Stem	0		0	

In the Table (b) the CAT activity in Lbc₃-5'-3'-CAT transformed L5-9 and L6-23 plants has been stated as the amount of ^{14}C -chloroamphenicol converted into acetylated derivatives. The amount of radio-activity in the acetylated derivatives has been

counted by liquid scintillation and stated in cpm/ μ g protein \cdot hour.

Example 6

Transcription test (Northern analysis) on tissues
5 of Lbc₃-5'-3'-CAT transformed and Lbc₃ transformed
Lotus plant lines.




5 μ g of total RNA extracted from root (R), nodule (N) or leaves + stem (LS) and separated in formaldehyde agarose gels were transferred onto nitrocellulose. Column 1 contains 5 μ g of total RNA from 5 20-day-old soybean nodules as control plants. The columns 2-4 and 5-7 contain total RNA from root, nodule or leaves + stem, respectively, of the Lbc₃-5'-3'-CAT transformed lines L5-9 and L6-23. The columns 8-10 contain RNA from corresponding tissues 10 of bird's-foot trefoil transformed by means of A. rhizogenes carrying the Lbc₃ gene in the T_L-DNA. In (a) radioactive DNA of the CAT coding sequence has been used as a probe for hybridization. The organ-specific transcription of the Lbc₃-5'-3'-15 CAT gene in root nodules from the L5-9 and L6-23 lines appears from columns 3 and 6. In (b) the transcript for the constitutive ubiquitine gene(s) is visualized using a cDNA probe for the human ubiquitine gene for the hybridization. In (c) the 20 nodule-specific transcription of bird's-foot trefoil own leghemoglobin genes is shown. A cDNA probe of the Lba gene of soybean has been used for this hybridization.

Example 7

Determination of the transcription initiation site (CAP site) of the Lbc₃ promoter of soybean in transformed root nodules of bird's-foot trefoil.

	Size marker		
+	16-23	polyA+mRNA	
+	15-9	polyA+mRNA	
+	Control	polyA+mRNA	



—TATAAATAAGTATTGATGTCGAAGTTGTTCTAACT— / —AAATGAG

The position of the "CAP site" was determined on the nucleotide level by means of primer extension. A synthetic oligonucleotide 5'CAACGGTGGTATATCCAGTG3' complementary to the nucleotides 15-34 in the coding
5 sequence of the CAT gene was used as primer for the enzyme reverse transcriptase. As a result single-stranded cDNA was formed the length of which corresponds to the distance between the 5' end of the primer and the 5' end of the primed mRNA. A 83
10 nucleotide cDNA strand would be expected according to the knowledge of the transcription initiation site of soybean Lbc₃ gene. Columns 2, 3, and 4 from left to right show the produced DNA strands when the primer extension has been operated on
15 polyA⁺-purified mRNA from transformed root nodules of bird's-foot trefoil, transformed leaves + stem of bird's-foot trefoil, and untransformed root nodules of bird's-foot trefoil, respectively. The 85, 86, 87, 88, and 90 nucleotides long cDNA strand
20 shown in column 2 proved correctly Lbc₃ promoter function in bird's-foot trefoil. The CAP sites corresponding to the cDNA sequences generated are indicated with asterisks (*) on the partial sequence of the Lbc₃ 5'3'-CAT region given. In the
25 sequence the TATA box of the Lbc₃ promoter and the corresponding translation initiation codon of the CAT coding sequence are underlined.

Example 8

Demonstration of the correct developmental control of the Lbc₃-5'-3'-CAT gene in transformed plants of bird's-foot trefoil (L6-23).

	Stage 1: No visible nodules	Stage 2: Emerging nodules	Stage 3: Distinct white nodules	Stage 4: Small pink nodules	Stage 5: Later stages of maturity
5 CAT activity					
in cpm/ μ g protein · hour	0	0	32.6	342.3	1255*
Nitrogenase activity					
nmol ethylene/ μ g protein · hour	0	0	0	0.5	2.7


10 * Substrate limited reaction; actual activity about 68000 cpm/ μ g protein · hour.

Chloroamphenicol acetyl transferase and nitrogenase activity were measured on cut off pieces of root with nodules at the different developmental stages indicated. The CAT activity can be detected in the white distinct nodules whereas the nitrogenase activity did not appear until the small pink nodules have developed. The latter development corresponds to the development known from soybean control plants and described by Marcker et al. EMBO J. 1984, 3, 1691-95. The CAT activity was determined as in Example 5. The nitrogenase activity was measured

as acetylene reduction capacity of the nodules followed by gaschromatographic determination of ethylene.

Example 9

- 5 Demonstration of Lbc₃ protein in bird's-foot trefoil plants transformed with the soybean Lbc₃ gene.




Proteins extracted from root nodules of Lbc₃ transformed (L8-35), Lbc₃-5'-3'-CAT transformed and nontransformed plants were separated by isoelectric focussing at a pH gradient of 4 to 5. The columns 1, 3, 5, 7, and 9 show Lbc₁, Lbc₂, Lbc₃, and Lba proteins synthesized in soybean control root nodules. Column 2 shows proteins from root nodules of Lbc₃-5'-3'-CAT transformed L6-23-bird's-foot trefoil plants, whereas the columns 6 and 8 show proteins from nontransformed plants. The columns 4 and 10 show soybean Lbc₃ protein synthesized in root nod-

ules of bird's-foot trefoil plants (L8-35) transformed with the Lbc₃ gene. The Lbc₃ protein band is indicated by an arrow.

Example 10

5 Expression of the Lbc₃-5'-3'-CAT gene requires the 5' Lbc₃ promoter region.

The Lbc₃-5'-3'-CAT gene construction carries a 2 Kb 5' Lbc₃ promoter region. Stepwise removal of sequences from the 5' end of this region demonstrated
10 that this promoter region is required for the characteristic expression of the Lbc₃-5'3'-CAT gene.



The Lbc₃-5'-3'-CAT gene construction was opened in
15 the unique XbaI site shown above, and digested with the exonuclease Bal31. A SalI linker fragment was ligated onto the blunt ends generated and the shortened SalI fragments carrying the Lbc₃-5'-3'-CAT gene were transferred into L.corniculatus. The effect
20 of removing promoter sequences was measured as CAT activity. End points of the deleted 5' region are given as the distance from the CAP site in nucleotides.

		CAT activity Cpm/ μ g protein/hrs.		
5' Lbc ₃	3' Lbc ₃	Root	Nodule	Leaf
2000	CAT	0	80000	0
-950		0	10000	0
-474		0	3000	0
-230		0	3000	0
-78		0	0	0

5 The drastically reduced level of CAT activity expressed from the Lbc₃ promoter deleted to nucleotide -230 and the zero activity from the promoter deleted to nucleotide -78 demonstrates that the Lbc₃ promoter region is required for the root nodule specific expression of the Lbc₃-5'-3'-CAT gene.


Example 11

Construction of the N23-CAT gene.

The N23 gene was isolated from a soybean DNA library as described in the enclosed paper of Sandal, Bojsen
15 and Marcker. The N23-CAT gene was constructed from the modified Lbc₃-5'-3'-CAT gene carried on plasmid pEJ5'-3'-CAT101 as described in the Applicant's
20 copending application No. 86 11 4704.9 concerning "Expression of Genes in Yeast", and a 1 Kb. EcoRI, DdeI fragment containing the N23 5' promoter region. The position of the EcoRI and DdeI sites in the N23 promoter region is indicated on the DNA sequence shown below. The cloning procedure used is outlined

below. The disclosure of the papers of Sandal et al., the EP application, and the paper of Jensen et al., Nature 321 (12 June 1986), 669-674, including the references cited should be considered incorporated into the present description as a means to amend, illustrate, and clarify it.

The N23-CAT gene was transferred to plants by the same method as the Lbc₃-5'-3'-CAT gene.



0249676

58

DNA sequence of the 5'-promotor region from the
N23 gene

```
      10      20      30      40      50      60      70
GAATTCGAGCTCGCCCGGGGATCGATCCTCTAGAGTCGACCTGCAGCCCAAGCTTGGATCAATCAATTAA
EcoRI                               SalI
5  TTCTATTGAGACACGATTGGAACAATTTTACATTATGAGACTATTTTGGTTTTTTTATTTGATCCAAA

      150     160     170     180     190     200     210
AAATTTAAAGCTTTAGATGATGATGAATTGAANNAATATTGTATTAATNTGAAAAGTTNNNNNGGTTTA

      220     230     240     250     260     270     280
ATGAATGCTATGATATTGATGGTCTTGATNTATTNNCAGAATTGAAAGTATTAAGAGAAGTGTTAAGAAA

10  AGAAGTTAGCACACCAATAGAAAGTATTGAGTTATATTAAACTTTAGATTCTTTTCAAATGTTTACATTG

      360     370     380     390     400     410     420
CATATAGAATTTTATTGACAATCCTTATAACAGTTGCTACTGTTGAAAGACGTTCTTCAAATTTAAATTT

      430     440     450     460     470     480     490
ACTTAAATCATATCTAAATCAACAATGTTACAAGATAGATTGAATGAGTTAGTTATTTTATCTATTGAA

15  AGTAAAGTGTTAGAATTGTTTGATTATAAACTCTGATAAATGATTTTGCAAGTTAAAAAACTAGAAGAT

      570     580     590     600     610     620     630
TAATATAAAAATTGATATTTTATATAATATATTAAGTCTCTTTAAATTCCTTGTAAGAAAAAGACATTTTT

      640     650     660     670     680     690     700
AAATAATAAATAAAGCAACTCTTAATTTTAATGAAACATCCCTTTGTTAAACCGAATCTTCATAATGT

20  AAAAATTAATGCTTGATGGAAGTTTTAATTTGTTCTACTCAATACTCAAAGGGTTGTAAATATTTTTTT

      780     790     800     810     820     830     840
TATCATTATATGTTGTAAATATGAATGCACTAGTAATTAGTTTAAATGATAAAATATATTCTACAGATAT

      850     860     870     880     890     900     910
ATTTCTGTCTCTTGGCAACTCGTGAGAATTGAATATATTATAAAGATGAAAGGTCGTTACAATTTTTTTT

25  AGAATAAATATTTATATACAATTCCTAGATTTTGTATATAAATTCACATATTGTATGAGTATAAATACAT

      990     1000     1010     1020     1030     1040     1050
GAGCACACACCAAACCTAGTCTCAAATTAAGTAAGGTGCTAATTATTAGCGGCTAGCTAAGTAACCAAGTA
DdeI
ATTAATG
```

Example 12

Organ-specific expression of the soybean N23-CAT gene in root nodules of *L.corniculatus* and *Trifolium repens*.

5 The activity of chloroamphenicol acetyl transferase (CAT) was measured as in example 5 and is given in cpm/ μ g protein/hrs.

10	<u>Table a.</u>		CAT activity	
	N23-CAT transformed <u><i>L.corniculatus</i></u>		Untransformed <u><i>L.corniculatus</i></u>	
	Root nodule	86150		0
	Root	0		0

15	<u>Table b.</u>		CAT activity	
	N23-CAT transformed <u><i>T.repens</i></u>		Untransformed <u><i>T.repens</i></u>	
	Root nodule	148000		0
	Root	0		0

20 Table (a) and b) shows the organ-specific expression of the N23-CAT gene in root nodules of *L.corniculatus* and *T.repens*. *L.corniculatus* was inoculated with *Rhizobium loti*, while *T.repens* was inoculated with *Rhizobium trifolii*.

In connection with the invention it has thus been proved that root nodule-specific genes can be expressed organ-specifically upon transfer to other
25 plants, here *Lotus corniculatus* and *Trifolium re-*

pens. It has furthermore been proved that the 5' flanking regions comprising the promoter are controlled by the organ-specific regulatory mechanism as the organ-specific control of the Lbc₃-5'-3'-CAT gene in Lotus corniculatus took place at the transcription level. The Lbc₃-5'-3'-CAT gene transferred was thus only transcribed in root nodules of transformed plants and not in other organs such as roots, stems, and leaves.

- 10 The expression of the Lbc₃-5'-3'-CAT gene in root nodules of transformed plants also followed the developmental timing known from soybean root nodules. No CAT activity could be detected in roots or small white root nodules (Example 8). A low
15 activity was present in the further developed white distinct nodules, whereas a high activity could be measured in the small pink nodules and mature nodules developed later on.

The organ-specific expression and the correct developmental expression of transferred root nodule-specific genes, here exemplified by the Lbc₃-5'-3'-CAT gene, allows as a particular use a functional expression of root nodule-specific genes also in other plants beyond leguminous plants. When all
20 the root nodule-specific plant genes necessary for the formation of root nodules are transferred from a leguminous plant to a non-root-nodule-forming plant species, the correct organ-specific expression proved above allows production of functionally
25 active, nitrogen-fixing root nodules on this plant upon infection by Rhizobium. In this manner these plants can grow without the supply of external

inorganic or organic nitrogen compounds. Root nodule-specific promoters, here exemplified by the Lbc₃ and N23 promoters, must be used in the present case for regulating the expression of the transferred genes.

According to the present invention a root nodule-specific promoter is used for expressing genes. The gene product or function of the gene product improves the function of the root nodule, e.g. by altering the oxygen transport, the metabolism, the nitrogen fixation or the nitrogen absorption.

Root nodules are thus used for the synthesis of biological products improving the plant per se or which can be extracted from the plant later on. A root nodule-specific promoter can be used for expressing a gene. The gene product or compound formed by said gene product constitute the desired product(s).

In connection with the present invention it has furthermore been proved that the soybean Lbc₃ leg-hemoglobin protein per se, i.e. the Lbc₃ gene product, is present in a high concentration in root nodules of bird's-foot trefoil plants expressing the Lbc₃ code sequence under the control of the Lbc₃ promoter. The latter has been proved by cloning the genomic Lbc₃ gene of the soybean into the integration vector pAR1, said genomic Lbc₃ gene containing the coding sequence, the intervening sequences, and the 5' and 3' flanking sequences. A 3.6 Kb BamHI fragment Lbc₃HH, cf. Example 2, was cloned into the pAR1 plasmid and transferred to

bird's-foot trefoil as stated previously.

The high level of Lbc₃ protein, cf. Example 9, found in transformed root nodules of bird's-foot trefoil and corresponding to the level in soybean root nodules proves an efficient transcription of the Lbc₃ promoter and an efficient processing and translation of Lbc₃mRNA in bird's-foot trefoil.

The high level of the CAT activity present in transformed root nodules is also a result of an efficient translation of mRNA formed from the chimeric Lbc₃ gene. The leader sequence on the Lbc₃ gene is decisive for the translation initiation and must determine the final translation efficiency. This efficiency is of importance for an efficient synthesis of gene products in plants or plant cells. An Lbc₃ or another leghemoglobin leader sequence can thus be used for increasing the final expression level of a predetermined plant promoter. The construction of a DNA fragment comprising a Lb leader sequence as first sequence and an arbitrary promoter as second sequence is a particular use of the invention when the construction is transferred and expressed in plants.

During nodule development around 30 different plant encoded polypeptides (nodulins) are specifically synthesized. Apart from the leghemoglobins, nodulins include nodule-specific forms of uricase (Bergmann et al (1983) EMBO. J. 2, 2333-2339), glutamine synthetase (Cullimore et al (1984) J.Mol. Appl. Genetics 2, 589-599) and sucrose synthase (Morell and Copeland (1985) Plant. Physiol. 78,

149-154). The function of most nodulins are, however, at present unknown.

Many nodulin genes have nevertheless been isolated and characterised during the last five years. These 5 include nodulins from several different legumes. Examples of such isolations and characterisations are widespread in the literature such as (Fuller et al (1983) Proc. Natl. Acad.Sci. 80, 2594-2598), (Sengupta-Gopalan et al (1986) Molec. Gen. Genet. 10 203, 410-420), (Bisseling et al (1985) in Proceedings of the 6th Int. symp. on Nitrogen Fixation, Martinus Nijhoff Publishers pp 53-59.), and (Gebhardt et al (1986) EMBO.J.5, 1429-1435). All of these genes contain nodule-specific regulatory 15 sequences. Such sequences and in fact entire 5' flanking regions and 3' flanking regions can furthermore be synthesized by automated oligonucleotide synthesis knowing the DNA sequences for the Lbc3 and N23 genes given in this description. Entire 20 nodule-specific genes can also be isolated with known recombinant techniques as described in the above papers and by (Maniatis et al (1982) Molecular cloning. A Laboratory Manual, Cold Spring Harbour Laboratory, New York).

25 The described method to obtain nodule-specific expression of genes can thus be reconstructed and performed according to the invention by any one skilled in the art of molecular genetics.

The method to obtain nodule-specific expression is 30 not dependent on the A. rhizogenes plant transformation described. Any other plant transformation

system e.g. A. tumefaciens systems, direct gene transfer or microinjection can equally be applied.

The A. rhizogenes system has been used and characterised by a number of scientific groups and is thus well-known from the literature. The characteristics of the system is described in:

Willmitzer et al. (1982), Molec.Gen.
Genet. 186, 16-22,

Chilton et al. (1982), Nature 295, 432-434,

10 Simpson et al. (1986), Plant.Molec.Biol.
6, 493-415,

Tepfer D. (1983), Molecular Genetics of
the Bacteria - Plant interaction,

15 Springer Verlag, Berlin Heidelberg pp
248-258,

White and Nester (1980), J.Bact. 144,
710-720,

Jaynes and Strobel (1981), Int.Rev. of Cytol.
Sup. 13, 105-125,

20 White and Nester (1980), J. Bact. 141,
1134-1141,

Pomponi et al. (1983), Plasmid 10, 119-
129, and

Slightom et al. (1986), J. Biol. Chem.
261, 108-121.

The latter two publications describe the restriction map and nucleotide sequence of the A. rhizogenes 5T_L-DNA segment used in the transformation system described here. With this information it is possible to anybody skilled in molecular genetics to use and reconstruct the "intermediate vectors" and the A. rhizogenes strains described here.

Claims:

1. A method of expressing genes in plants, parts of plants, and plant cell cultures by introducing into a cell thereof a recombinant DNA segment containing both the gene to be expressed and a 5' flanking region comprising a promoter sequence, and optionally a 3' flanking region, and culturing of the transformed cells in a growth medium, characterised by using as the recombinant DNA segment a DNA fragment comprising an inducible plant promoter (as defined) from root nodule-specific genes.
2. A method as claimed in claim 1, characterised by using a DNA fragment comprising an inducible plant promoter (as defined) and being identical with, derived from or comprising 5' flanking regions of root nodule-specific genes.
3. A method as claimed in claim 2, characterised by using a DNA fragment comprising an inducible plant promoter (as defined) and being identical with, derived from or comprising 5' flanking regions of root nodule-specific genes, said DNA fragment causing an expression of a gene which is induced in root nodules at specific stages of development and as a step of the symbiosis, whereby nitrogen fixation occurs.
4. A method as claimed in claims 1-3 for the expression of root nodule-specific genes, characterised by using a DNA fragment comprising an inducible plant promoter (as defined)

from root nodule-specific genes.

5. A method as claimed in claims 1-3 for the expression of genes in leguminous plants, parts of leguminous plants, and leguminous plant cell cultures, characterised by using a DNA fragment comprising an inducible plant promoter (as defined) from root nodule-specific genes.

6. A method as claimed in claims 1-5, characterised by the DNA fragment comprising the inducible plant promoter and being identical with, derived from or comprising 5' flanking regions of leghemoglobin genes.

7. A method as claimed in claim 6, characterised by the DNA fragment comprising the inducible plant promoter and being identical with, derived from or comprising 5' flanking regions of soybean leghemoglobin genes.

8. A method as claimed in claim 7, characterised by the DNA fragment comprising the inducible plant promoter and being identical with, derived from or comprising 5' flanking regions of the Lba gene with the sequence

```

25 GAGATACATT ATAATAATCT CTCTAGTGTC TATTTATTAT TTTATCTGGT
   GATATATACC TTCTCGTATA CTGTTATTTT TTCAATCTTG TAGATTTACT
   TCTTTTATTT TTATAAAAAA GACTTTATTT TTTTAAAAAA AATAAAGTGA
   ATTTTGAAAA CATGCTCTTT GACAATTTTC TGTTTCCTTT TTCATCATTG
   GGTAAATCT CATAGTGCCT CTATTCAATA ATTTGGGCTC AATTTAATTA
   GTAGAGTCTA CATAAAATTT ACCTTAATAG TAGAGAATAG AGAGTCTTGG
   AAAGTTGGTT TTTCTCGAGG AAGAAAGGAA ATGTTAAAAA CTGTGATATT
   TTTTTTTTGG ATTAATAGTT ATGTTTATAT GAAAACTGAA AATAAATAAA
   CTAACCATAT TAAATTTAGA ACAACACTTC AATTATTTTT TTAATTTGAT
   TAATTAAAAA ATTATTTGAT TAAATTTTTT AAAAGATCGT TGTTTCTTCT
   TCATCATGCT GATTGACACC CTCCACAAGC CAAGAGAAAC ACATAAGCTT
30 TGGTTTTCTC ACTCTCCAAG CCCTCTATAT AAACAAATAT TGGAGTGAAG

```

TTGTTGCATA ACTTGCATCG AACAAATTAAT AGAAATAACA GAAATTTAAA
AAAGAAATAT G.

9. A method as claimed in claim 7, c h a r -
a c t e r i s e d by the DNA fragment comprising
5 the inducible plant promoter and being identical
with, derived from or comprising 5' flanking regions
of the Lbc₁ gene with the sequence:

10 TTCTCTTAAT ACAATGGAGT TTTTGTGAA CATAATACA TTAAAAAAA
AATCTCTAGT GTCTATTTAC CCGGTGAGAA GCCTTCTCGT GTTTTACACA
CTTTAATATT ATTATATCCT CAACCCACACA AAAAAGAATA CTGTTATATC
TTTCCAAACC TGTAGATTTA TTTATTTATT TATTTATTTT TACAAAGGAG
ACTTCAGAAA AGTAATTACA TAAAGATAGT GAACATCATT TTATTTATTA
TAATAAACTT TAAAATCAAA CTTTTTTTATA TTTTTTGTGA CCCTTTTTCAT
TATTGGGTGA AATCTCATAG TGAAGCCATT AAATAATTTG GGCTCAAGTT
TTATTAGTAA AGTCTGCATG AAATTTAACT TAACAATAGA GAGAGTTTTC
15 GAAAGGGAGC GAATGTTAAA AAGTGTGATA TTATATTTTA TTTCGATTAA
TAATTATGTT TACATGAAAA CATACAAAAA AATACTTTTA AATTCAGAAT
AATACTTAAA ATATTTATTT GCTTAATTGA TTAAGTGAAT ATTATTTGAT
TAGGATTTTG AAAAGATCAT TGGCTCTTCG TCATGCCGAT TGACACCCCTC
CACAAGCCAA GAGAACTTA AGTTGTAAAC TTTCTCACTC CAAGCCTTCT
ATATAAACAT GTATTGGATG TGAAGTTATT GCATAACTTG CATTGAACAA
TAGAAAAATA CAAAAAAAG TAAAAAAGTA GAAAAGAAAT ATG,

- 20 10. A method as claimed in claim 7, c h a r -
a c t e r i s e d by the DNA fragment comprising
the inducible plant promoter and being identical
with, derived from or comprising 5' flanking regions
of the Lbc₂ gene with the sequence:

25 TCGAGTTTTT ACTGAACATA CATTTATTAA AAAAAACTCT CTAGTGTCCA
TTTATTCGGC GAGAAGCCTT CTCGTGCTTT ACACACTTTA ATATTATTAT
ATCCCCACCC CCACCAAAAA AAAAAAAACT GTTATATCTT TCCAGTACAT
TTATTTCTTA TTTTACAAA GGAAACTTCA CGAAAGTAAT TACAAAAAAG
ATAGTGAACA TCATTTTTTT AGTTAAGATG AATTTTAAAA TCACACTTTT
TTATATTTT TTGTTACCCT TTTCATTATT GGGTGAAATC TCATAGTGAA
ACTATTAAAT AGTTTGGGCT CAAGTTTAT TAGTAAAGTC TGCATGAAAT
TTAACTTAAT AATAGAGAGA GTTTTGGAAA GGTAACGAAT GTTAGAAAGT
30 GTGATATTAT TATAGTTTAA TTTAGATTAA TAATTATGTT TACATGAAAA
TTGACAATTT ATTTTAAAA TTCAGAGTAA TACTTAAATT ACTTATTTAC
TTTAAGATTT TGAAAAGATC ATTTGGCTCT TCATCATGCC GATTGACACC
CTCCACAAGC CAAGAGAAAC TTAAGTTGTA ATTTTCTTAA CTCCAGCCT
TCTATATAAA CACGTATTGG ATGTGAAGTT GTTGCATAAC TTGCATTGAA
CAATAGAAAT AACAAACAAAG AAAATAAGTG AAAAAAGAAA TATG,

11. A method as claimed in claim 7, c h a r -
a c t e r i s e d by the DNA fragment comprising
the inducible plant promoter and being identical
with, derived from or comprising 5' flanking regions
5 of the Lbc₃ gene with the sequence:

```

TATGAAGATT AAAAAATACA CTCATATATA TGCCATAAGA ACCAACAAAA
GTACTATTTA AGAAAAGAAA AAAAAAACCT GCTACATAAT TTCCAATCCTT
GTAGATTTAT TTCTTTTATT TTTATAAAGG AGAGTTAAAA AAATTACAAA
ATAAAAATAG TGAACATCGT CTAAGCATTT TTATATAAGA TGAATTTTAA
10 AAATATAATT TTTTGTGCTA AATCGTATGT ATCTTGTCTT AGAGCCATTT
TTGTTTAAAT TGGATAAGAT CACACTATAA AGTTCTTCCT CCGAGTTTGA
TATAAAAAAA ATTGTTTCCC TTTTGATTAT TGGATAAAAT CTCGTAGTGA
CATTATATTA AAAAAATTAG GGCTCAATTT TTATTAGTAT AGTTTGCATA
AATTTTAACT TAAAAATAGA GAAAATCTGG AAAAGGGACT GTTAAAAAGT
GTGATATTAG AAATTTGTCG GATATATTAA TATTTTATTT TATATGGAAA
CTAAAAAAAT ATATATTAAA ATTTTAAATT CAGAATAATA CTTAAATTAT
TTATTTACTG AAAATGAGTT GATTTAAGTT TTTGAAAAGA TGATTGTCTC
15 TTCACCATAC CAATTGATCA CCCTCCTCCA ACAAGCCAAG AGAGACATAA
GTTTTATTAG TTATTCTGAT CACTCTTCAA GCCTTCTATA TAAATAAGTA
TTGGATGTGA AGTTGTTGCA TAACTTGCAT TGAACAATTA ATAGAAATAA
CAGAAAAGTA GAAAAGAAAT ATG.

```

12. A method as claimed in claim 7, c h a r a c -
20 t e r i s e d by the DNA fragment comprising the
inducible plant promoter and being identical with,
derived from or comprising 5' flanking regions of
the Lbc₃-5'-3'-CAT gene with the sequence:

```

TATGAAGATT AAAAAATACA CTCATATATA TGCCATAAGA ACCAACAAAA
25 GTACTATTTA AGAAAAGAAA AAAAAAACCT GCTACATAAT TTCCAATCCTT
GTAGATTTAT TTCTTTTATT TTTATAAAGG AGAGTTAAAA AAATTACAAA
ATAAAAATAG TGAACATCGT CTAAGCATTT TTATATAAGA TGAATTTTAA
AAATATAATT TTTTGTGCTA AATCGTATGT ATCTTGTCTT AGAGCCATTT
TTGTTTAAAT TGGATAAGAT CACACTATAA AGTTCTTCCT CCGAGTTTGA
TATAAAAAAA ATTGTTTCCC TTTTGATTAT TGGATAAAAT CTCGTAGTGA
30 CATTATATTA AAAAAATTAG GGCTCAATTT TTATTAGTAT AGTTTGCATA
AATTTTAACT TAAAAATAGA GAAAATCTGG AAAAGGGACT GTTAAAAAGT
GTGATATTAG AAATTTGTCG GATATATTAA TATTTTATTT TATATGGAAA
CTAAAAAAAT ATATATTAAA ATTTTAAATT CAGAATAATA CTTAAATTAT
TTATTTACTG AAAATGAGTT GATTTAAGTT TTTGAAAAGA TGATTGTCTC
TTCACCATAC CAATTGATCA CCCTCCTCCA ACAAGCCAAG AGAGACATAA
GTTTTATTAG TTATTCTGAT CACTCTTCAA GCCTTCTATA TAAATAAGTA
TTGGATGTGA AGTTGTTGCA TAACTTGCAT TGAACAATTA ATAGAAATAA
CAGAAAAGTA GAATTCTAAA ATG

```

13. A method as claimed in claim 5, c h a r a c -
t e r i s e d by the DNA fragment comprising the
inducible plant promoter and being identical with,
derived from or comprising 5' flanking regions of
5 the N23 gene with the sequence:

```

      10      20      30      40      50      60      70
GAATTCGAGCTCGCCCGGGGATCGATCCTCTAGAGTCGACCTGCAGCCCAAGCTTGGATCAATCAATTAA
EcoRI                               SalI
      80      90     100     110     120     130     140
TTCTATTGAGACACGATTTGAACAATTTTACATTATGAGACTATTTTGGTTTTTTATTGATCCAAAA

10  AAATTTAAAGCTTTAGATGATGATGAATTGAANNAATATTGTATTAAATNNTGAAAAGTTNNNNNGGTTTA

      220     230     240     250     260     270     280
ATGAATGCTATGATATTGATGGTCTTGATNTATTNNCAGAATTGAAAGTATTAAGAGAAGTCTTAAGAAA

      290     300     310     320     330     340     350
AGAAGTTAGCACACCAATAGAAAGTATTGAGTTATATTAAACTTTAGATTCTTTTCAATGTTTACATTG

15  CATATAGAATTTTATTGACAATCCTTATAACAGTTGCTACTGTTGAAAGACGTTCTTCAAATTTAAATTT

      430     440     450     460     470     480     490
ACTTAAATCATATCTAAAATCAACAATGTTACAAGATAGATTGAATGAGTTAGTTATTTTATCTATTGAA

      500     510     520     530     540     550     560
AGTAAAGTGTTAGAATTGTTTGATTATAAACTCTGATAAATGATTTTGCAGTTAAAAAACTAGAAAGAT

20  TAATATAAAAATTGATATTTTATATAATATATTAAAGTCTCTTTAAAATTCTTGTAaaaaaAGACATTTT

      640     650     660     670     680     690     700
AAATAATAAAATAAAGCAACTCTTAATTTTAATGAAACATCCCTTTGTTAAACCGAATCTTCCATAATGT

      710     720     730     740     750     760     770
AAAAATTAATGCTTGATGGAAGTTTTTAATTTGTTCTACTCAATACTCAAAGGGTTGTAATATTTTTTTT

25  TATCATTATATGTTGTAAATATGAATGCACTAGTAATTAGTTTAAATGATAAAATATATTCTACAGATAT

      850     860     870     880     890     900     910
ATTTCTGTCTCTTGGCAACTCGTGAGAATTGAATATATTATAAAGATGAAAGGTCGTTACAATTTTTTTT

      920     930     940     950     960     970     980
AGAATAAATATTTATATACAATTCCTAGATTTTGTATAAAATTCACATATTGTATGAGTATAAATACAT

30  GAGCACACACCAAACCTAGTCTCAAATTAAGTAAGGTGCTAATTATTAGCGGCTAGCTAAGTAACCAAGTA
      DdeI

      ATTAATG
```

14. A method as claimed in any of the claims 1-13, characterised by the 3' flanking region of the genes to be expressed being a 3' flanking region of root nodule-specific genes of any origin.

15. A method as claimed in claim 14, characterised by the 3' flanking region being of leghemoglobin genes.

16. A method as claimed in claim 14, characterised by the 3' flanking region being of soybean leghemoglobin genes.

17. A method as claimed in claim 16, characterised by the 3' flanking region being of the Lba, Lbc₁, Lbc₂ or Lbc₃ gene with the following sequences, respectively:

Lba

1590
TAA TTA GTA TCT ATT GCA GTA AAG TGT AAT AAA TAA ATC TTG 1620

1650
20 TTT CAC TAT AAA ACT TGT TAC TAT TAG ACA AGG GCC TGA TAC AAA ATG TTG GTT AAA ATA 1680

1710
ATG GAA TTA TAT AGT ATT GGA TAA AAA TCT TAA GGT TAA TAT TCT ATA TTT GCG TAG GTT 1740

1770
TAT GCT TGT GAA TCA TTA TCG GTA TTT TTT TTC CTT TCT GAT AAT TAA TCG GTA AAT TA 1800

1830
25 ACA AAT AAG TTC AAA ATG ATT TAT ATG TTT CAA AAT TAT TTT AAC AGC AGG TAA AAT GTT 1860

ATT TGG TAC GAA AGC TAA TTC GTC GA

0249676

72

Lbc₁

1320
TAA/TT AGG ATC TAC TGC ATT GCC GTA

1350 1380
AAG TGT AAT AAA TAA ATC TTG TTT CAA CTA AAA CTT GTT ATT AAA CAA GTT CCC TAT ATA

1410 1440
AAT GTT GTT TAA AAT AAG TAA ATT TCA TTG TAT TGG ATA AAC ACT TTT AAG TTA TAT ATT

5 1470 1500
TCC ATA TAT TTA CGT TTG TGA ATC ATA ATC GAT ACT TTA TAA AAA TAA ATT CCA AAT AAT

TTA TAC GTT TTA AAA ATT ATT TT

Lbc₂

10

TAG/GAT CTA CTA TTG CCG TCA AGT
1140

GTA ATA AAT AAA TTT TGT TTC ACT AAA ACT TGT TAT TAA ACA AGT CCC CGA TAT ATA AAT
1170 1200

GTT GGT TAA AAT AAG TAA ATT ATA CGG TAT TGA TAA ACA ATC TTA AGT TTT ATA TAT AGT
1230 1260

TCC ATA TAC TAA AGT TTG TGA ATC ATA ATC GA
1290

15 and Lbc₃

TAG/GAT CTA CAA TTG CCT TAA AGT GTA ATA AAT AAA
990 1020

TAT TAT TTC ACT AAA ACT TGT TAT TAA ACC AAG TTC TCG ATA TAA ATG TTG GTT AAA CTA
1050 1080

20 AGT AAA TTA TAT GGT ATT GGA TAA ACA ATC TTA AGC TT
1110

18. A method as claimed in claim 1 of preparing
a polypeptide by introducing into a cell of a plant,
a part of a plant or a plant cell culture a recombi-
25 nant plasmid, c h a r a c t e r i s e d by using
as the recombinant plasmid a plasmid comprising an
inducible plant promoter (as defined) of root nod-
ule-specific genes.

19. A DNA fragment comprising an inducible plant promoter (as defined) to be used when carrying out the method as claimed in claims 1-18, c h a r -
a c t e r i s e d by being identical with, de-
5 rived from or comprising a 5' flanking region of root nodule-specific genes of any origin.

20. A DNA fragment as claimed in claim 19,
c h a r a c t e r i s e d by being identical with,
derived from or comprising a 5' flanking region of
10 plant leghemoglobin genes.

21. A DNA fragment as claimed in claim 20,
c h a r a c t e r i s e d by being identical with,
derived from or comprising a 5' flanking region of
soybean leghemoglobin genes.

15 22. A DNA fragment as claimed in claim 21,
c h a r a c t e r i s e d by being identical with,
derived from or comprising a 5' flanking region of
the Lba gene with the sequence:

	GAGATACATT	ATAATAATCT	CTCTAGTGTC	TATTTATTAT	TTTATCTGGT
20	GATATATACC	TTCTCGTATA	CTGTTATTTT	TTCAATCTTG	TAGATTTACT
	TCTTTTATTT	TTATAAAAAA	GACTTTATTT	TTTTAAAAAA	AATAAAGTGA
	ATTTTGAAAA	CATGCTCTTT	GACAATTTTC	TGTTTCCTTT	TTCATCATTG
	GGTTAAATCT	CATAGTGCCT	CTATTCAATA	ATTTGGGCTC	AATTTAATTA
	GTAGAGTCTA	CATAAAATTT	ACCTTAATAG	TAGAGAATAG	AGAGTCTTGG
	AAAGTTGGTT	TTTCTCGAGG	AAGAAAGGAA	ATGTTAAAAA	CTGTGATATT
	TTTTTTTTTG	ATTAATAGTT	ATGTTTATAT	GAAAAC TGAA	AATAAATAAA
25	CTAACCATAT	TAAATTTAGA	ACAACACTTC	AATTATTTTT	TTAATTTGAT
	TAATTAAAAA	ATTATTTGAT	TAAATTTTTT	AAAAGATCGT	TGTTTCTTCT
	TCATCATGCT	GATTGACACC	CTCCACAAGC	CAAGAGAAAC	ACATAAGCTT
	TGGTTTTCTC	ACTCTCCAAG	CCCTCTATAT	AAACAAATAT	TGGAGTGAAG
	TTGTTGCATA	ACTTGCATCG	AACAATTAAT	AGAAATAACA	GAAAATTAAA
	AAAGAAATAT	G,			

23. A DNA fragment as claimed in claim 21,
c h a r a c t e r i s e d by being identical with,
derived from or comprising a 5' flanking region of
the Lbc₁ gene with the sequence:

```

TTCTCTTAAT ACAATGGAGT TTTTGTTGAA CATACTATACA TTTAAAAAAA
5 AATCTCTAGT GTCTATTTAC CCGGTGAGAA GCCTTCTCGT GTTTTACACA
CTTTAATATT ATTATATCCT CAACCCACACA AAAAAGAATA CTGTTATATC
TTTCCAAACC TGTAGATTTA TTTATTTATT TATTTATTTT TACAAAGGAG
ACTTCAGAAA AGTAATTACA TAAAGATAGT GAACATCATT TTATTTATTA
TAATAAACTT TAAAATCAAA CTTTTTTTATA TTTTTTGTTA CCCTTTTCAT
TATTGGGTGA AATCTCATAG TGAAGCCATT AAATAATTTG GGCTCAAGTT
TTATTAGTAA AGTCTGCATG AAATTTTAACT TAACAATAGA GAGAGTTTTC
10 GAAAGGGAGC GAATGTTAAA AAGTGTGATA TTATATTTTA TTTCGATTAA
TAATTATGTT TACATGAAAA CATACAAAAA AATACTTTTA AATTCAGAAT
AATACTTAAA ATATTTATTT GCTTAATTGA TTAAGTGAAT ATTATTTGAT
TAGGATTTTG AAAAGATCAT TGGCTCTTCG TCATGCCGAT TGACACCCCTC
CACAAAGCCAA GAGAAACTTA AGTTGTAAAC TTTCTCACTC CAAGCCTTCT
ATATAAACAT GTATTGGATG TGAAGTTATT GCATAACTTG CATTGAACAA
TAGAAAAATA CAAAAAAAAG TAAAAAAGTA GAAAAGAAAT ATG,

```

15 24. A DNA fragment as claimed in claim 21,
c h a r a c t e r i s e d by being identical with,
derived from or comprising a 5' flanking region of
the Lbc₂ gene with the sequence:

```

TCGAGTTTTT ACTGAACATA CATTTATTAA AAAAAACTCT CTAGTGTCCA
TTTATTCGGC GAGAAGCCTT CTCGTGCTTT ACACACTTTA ATATTATTAT
20 ATCCCCACCC CCACCAAAAA AAAAAAACT GTTATATCTT TCCAGTACAT
TTATTTCTTA TTTTACAAA GGAACTTCA CGAAAGTAAT TACAAAAAAG
ATAGTGAACA TCATTTTTTT AGTTAAGATG AATTTTAAAA TCACACTTTT
TTATATTTTT TTGTTACCCT TTTCATTATT GGGTGAAATC TCATAGTGAA
ACTATTAAAT AGTTTGGGCT CAAGTTTTAT TAGTAAAGTC TGCATGAAAT
TTAACTTAAT AATAGAGAGA GTTTTGGAAA GGTAACGAAT GTTAGAAAGT
GTGATATTAT TATAGTTTTA TTTAGATTAA TAATTATGTT TACATGAAAA
TTGACAATTT ATTTTAAAA TTCAGAGTAA TACTTAAATT ACTTATTTAC
25 TTTAAGATTT TGAAAAGATC ATTTGGCTCT TCATCATGCC GATTGACACC
CTCCACAAGC CAAGAGAAAC TTAAGTTGTA ATTTTCTAA CTCCAAGCCT
TCTATATAAA CACGTATTGG ATGTGAAGTT GTTGCATAAC TTGCATTGAA
CAATAGAAAT AACAACAAAG AAAATAAGTG AAAAAAGAAA TATG,

```

25. A DNA fragment as claimed in claim 21,
c h a r a c t e r i s e d by being identical with,
30 derived from or comprising a 5' flanking region of

the Lbc₃ gene with the sequence:

```

TATGAAGATT AAAAAATACA CTCATATATA TGCCATAAGA ACCAACA AAAA
GTACTATTTA AGAAAAGAAA AAAAAAACCT GCTACATAAT TTCCAATCCTT
GTAGATTTAT TTCTTTTATT TTTATAAAGG AGAGTTAAAA AAATTACAAA
5 ATAAAAATAG TGAACATCGT CTAAGCATTT TTATATAAGA TGAATTTTAA
AAATATAATT TTTTGTCTA AATCGTATGT ATCTTGTCTT AGAGCCATTT
TTGTTTAAAT TGGATAAGAT CACACTATAA AGTTCTTCCT CCGAGTTTGA
TATAAAAAAA ATTGTTTCCC TTTTGATTAT TGGATAAAAT CTCGTAGTGA
CATTATATTA AAAAAATTAG GGCTCAATTT TTATTAGTAT AGTTTGCATA
AATTTTAACT TAAAAATAGA GAAATCTGG AAAAGGGACT GTTAAAAAGT
GTGATATTAG AAATTTGTCTG GATATATTAA TATTTTATTT TATATGGAAA
10 CTAAAAAAAT ATATATTAAA ATTTTAAATT CAGAAATAA CTTAAATTAT
TTATTTACTG AAAATGAGTT GATTTAAGTT TTTGAAAAGA TGATTGTCTC
TTCACCATAC CAATTGATCA CCCTCCTCCA ACAAGCCAAG AGAGACATAA
GTTTTATTAG TTATTCTGAT CACTCTTCAA GCCTTCTATA TAAATAAGTA
TTGGATGTGA AGTTGTTGCA TAACTTGCAT TGAACAATTA ATAGAAATAA
CAGAAAAGTA GAAAAGAAAT ATG.

```

- 15 26. A DNA fragment as claimed in claim 21,
c h a r a c t e r i s e d by the DNA fragment
comprising the inducible plant promoter being iden-
tical with, derived from or comprising 5' flanking
regions of Lbc₃-5'-3'-CAT gene with the sequence:

```

20 TATGAAGATT AAAAAATACA CTCATATATA TGCCATAAGA ACCAACA AAAA
GTACTATTTA AGAAAAGAAA AAAAAAACCT GCTACATAAT TTCCAATCCTT
GTAGATTTAT TTCTTTTATT TTTATAAAGG AGAGTTAAAA AAATTACAAA
ATAAAAAATAG TGAACATCGT CTAAGCATTT TTATATAAGA TGAATTTTAA
AAATATAATT TTTTGTCTA AATCGTATGT ATCTTGTCTT AGAGCCATTT
TTGTTTAAAT TGGATAAGAT CACACTATAA AGTTCTTCCT CCGAGTTTGA
TATAAAAAAA ATTGTTTCCC TTTTGATTAT TGGATAAAAT CTCGTAGTGA
25 CATTATATTA AAAAAATTAG GGCTCAATTT TTATTAGTAT AGTTTGCATA
AATTTTAACT TAAAAATAGA GAAATCTGG AAAAGGGACT GTTAAAAAGT
GTGATATTAG AAATTTGTCTG GATATATTAA TATTTTATTT TATATGGAAA
CTAAAAAAAT ATATATTAAA ATTTTAAATT CAGAAATAA CTTAAATTAT
TTATTTACTG AAAATGAGTT GATTTAAGTT TTTGAAAAGA TGATTGTCTC
TTCACCATAC CAATTGATCA CCCTCCTCCA ACAAGCCAAG AGAGACATAA
GTTTTATTAG TTATTCTGAT CACTCTTCAA GCCTTCTATA TAAATAAGTA
TTGGATGTGA AGTTGTTGCA TAACTTGCAT TGAACAATTA ATAGAAATAA
30 CAGAAAAGTA GAATTCTAAA ATG

```

27. A DNA fragment as claimed in claim 19,
c h a r a c t e r i s e d by being identical with,

derived from or comprising 5' flanking regions of the N23 gene with the sequence:

```

      10      20      30      40      50      60      70
GAATTCGAGCTCGCCCGGGGATCGATCCTCTAGAGTCGACCTGCAGCCCAAGCTTGGATCAATCAATTAA
  ECORI           SalI
5  TTCTATTGAGACACGATTGAACAATTTTACATTATGAGACTATTTTGGTTTTTTATTTGATCCAAA
      80      90      100     110     120     130     140
AAATTTAAGCTTTAGATGATGAATTGAANNAATATTGTATTAATNTGAAAAGTTNNNNNGGTTTA
      150     160     170     180     190     200     210
ATGAATGCTATGATATTGATGGTCTTGATNTATTNNCAGAATTGAAAGTATTAAGAGAAGTGTTAAGAA
      220     230     240     250     260     270     280
10 AGAAGTTAGCACACCAATAGAAAGTATTGAGTTATATTAAGAACTTTAGATTCTTTTCAATGTTTACATTG
      290     300     310     320     330     340     350
CATATAGAAATTTATTGACAATCCTTATAACAGTTGCTACTGTTGAAAGACGTTCTTCAAATTAAGATT
      360     370     380     390     400     410     420
ACTTAAATCATATCTAAATCAACAATGTTACAAGATAGATTGAATGAGTTAGTTATTTTATCTATTGAA
      430     440     450     460     470     480     490
15 AGTAAAGTGTTAGAAATGTTTGATTATAAACTCTGATAAATGATTTGCAGTTAAAAAACTAGAAGAT
      500     510     520     530     540     550     560
TAATATAAAATGATATTTTATATAATATATTAAGTCTCTTTAAATTCCTTGTAAGAAAAGACATTTT
      570     580     590     600     610     620     630
AAATAATAAAATAAGCAACTCTTAATTTAATGAAACATCCCTTTGTTAAACCGAATCTTCATAATGT
      640     650     660     670     680     690     700
20 AAAAATTAATGCTTGATGGAAGTTTAAATTTGTTCTACTCAATACTCAAAGGGTTGTAAATATTTTTT
      710     720     730     740     750     760     770
TATCATTATATGTTGTAAATATGAATGCACTAGTAATTAGTTTAAATGATAAAATATATTCTACAGATAT
      780     790     800     810     820     830     840
ATTCTGTCTCTTGGCAACTCGTGAGAATTGAATATATTATAAAGATGAAAGGTCGTTACAATTTTTTT
      850     860     870     880     890     900     910
25 AGAATAAATATTTATATACAATTCCTAGATTTTGTATATAAATTCACATATTGTATGAGTATAAATACAT
      920     930     940     950     960     970     980
GAGCACACACCAACTAGTCTCAAATTAAGTAAGGTGCTAATTATTAGCGGCTAGCTAAGTAACCAAGTA
      990    1000    1010    1020    1030    1040    1050
                                   DdeI
ATTAAATG

```

28. A plasmid which can be used when carrying

out the method as claimed in claims 1-18,
c h a r a c t e r i s e d by comprising a DNA
fragment as claimed in any of the claims 19-27.

29. A plasmid as claimed in claim 28, c h a r -
5 a c t e r i s e d by being pAR29.

30. A plasmid as claimed in claim 28, c h a r -
a c t e r i s e d by being pAR30.

31. A plasmid as claimed in claim 28, c h a r -
a c t e r i s e d by being pAR11.

10 32. A plasmid as claimed in claim 28, c h a r -
a c t e r i s e d by being N23-CAT.

33. A transformant Agrobacterium rhizogenes 15834-
strain which can be used when carrying out the
method as claimed in any of the claims 1 to 18,
15 c h a r a c t e r i s e d by the bacterium strain
being transformed by a plasmid according to any of
the preceding claims 28 to 32.

34. A transformant Agrobacterium rhizogenes 15834-
strain which can be used when carrying out the
20 method as claimed in any of the claims 1 to 18,
c h a r a c t e r i s e d by the bacterium strain
being transformed by pAR29 and being named AR1127.

35. A transformant Agrobacterium rhizogenes 15834-
strain which can be used when carrying out the
25 method as claimed in any of the claims 1 to 18,
c h a r a c t e r i s e d by the bacterium strain
being transformed by pAR30 and being named AR1134.

36. A transformant Agrobacterium rhizogenes 15834-strain which can be used when carrying out the method as claimed in any of the claims 1 to 18, characterised by the bacterium strain
5 being transformed by pAR11 and being named AR1000.

37. A transformant Agrobacterium rhizogenes 15834-strain which can be used when carrying out the method as claimed in any of the claims 1 to 18, characterised by the bacterium strain
10 being transformed by N23-CAT and being named AR204-N23-CAT.

38. Plants, parts of plants and plant cells, particularly of the family Leguminosae, obtainable by transformation with a recombinant DNA segment,
15 fragment or plasmid according to any one of the claims 1 to 37.